

Stereotaxic Atlas of the Marmoset Brain

with Immunohistochemical Architecture and MR Images

Volume 1
Nissl-stained sections
and their diagrams





Shigeki Yuasa Katsuki Nakamura Shinichi Kohsaka

National Institute of Neuroscience National Center of Neurology and Psychiatry, Japan

Stereotaxic Atlas of the Marmoset Brain

with immunohistochemical architecture and MR images

Volume 1

Nissl-stained sections and their diagrams

Shigeki Yuasa M.D., Ph.D.

Director, Department of Ultrastructural Research

Katsuki Nakamura Ph.D.

Director, Department of Animal Models for Human Disease

Shinichi Kohsaka M.D., Ph.D. Director-General

ISBN:978-4-9905092-0-0

Preface

As president of the National Center of Neurology and Psychiatry, it gives me great pleasure to announce the publication of *Stereotaxic Atlas of the Marmoset Brain*.

The main role of the National Center of Neurology and Psychiatry is to contribute to the investigation and elucidation of mental disorders, neurological disorders, muscular diseases, and developmental disorders, and to provide our fellow citizens with highly advanced medical treatment, based on the results of our research.

In addition to animal models such as mice and rats, the importance of translational research using primates has been on the increase. In particular, in studies of mental and developmental disorders such as autism, it is essential to investigate abnormalities of higher brain function, including communication disorders. In such studies, the marmoset may be a very valuable as animal model, since it forms a family of mother, father and children living as a unit and communicating with each other using animal sounds. As an animal model, the marmoset has several advantages over other primates in that it is relatively easy to handle and feed, weights only about 500 g, and delivers two or three babies twice a year.

On the other hand, there have also been difficulties in using the marmoset as an animal model. We have not had an accurate atlas showing brain structures that have been fully examined, as in the case of mouse or rat, no test method that determines abnormal behavior correctly, and inadequate information on genes and proteins. However, it has been the accumulation of solid sober research in our National Institute of Neuroscience that has solved the problems of using marmoset in research work.

The first result of our studies is this publication of two volumes of Stereotaxic Atlas of the *Marmoset Brain*. We have illustrated the books giving researchers the information they need actually as you see it. We sincerely hope that this publication will help readers both in Japan and overseas achieve dramatic progress in their research into elucidating higher brain function disorders using marmoset.

In closing, I would like to express my sincere gratitude to Dr. Shigeki Yuasa and all the researchers in the National Institute of Neuroscience for their immense contributions.

Teruhiko Higuchi, President National Center of Neurology and Psychiatry Tokyo, Japan

March, 2010

Acknowledgements

We are very grateful to Ms. Yumiko Kamagata, Department of Ultrastructural Research for her skillful assistance in the histological work. We are indebted to Drs. Akihiro Izumi, Hidetoshi Ishibashi, Atsuko Saito, Osamu Yokoyama, Shinya Uchida and Koji Kuraoka, Department of Animal Models for Human Disease for their technical assistance. We also thank Megumi Koyanagawa, Keiko Otokuni and Arisa Ishikawa for their daily support. This work has been supported in part by the Research Grant (20B-10) for Nervous and Mental Disorders from the Ministry of Health, Labor and Welfare of Japan and by Strategic Research Program for Brain Science from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

Contents

1.	Intro	duction	
		General remarks.	
	1.2.	General construction of this atlas	2
2.	Mate	erials and Methods	
	2.1.	Histological procedures	3
		MRI	
		Stereotaxic coordinates	
3.	Inde	x of Abbreviations	5
4.	Table	es of Plates	9
5.	Plate	s: Nissl-stained sections and corresponding tracing	11
6	Liter	ratures on the Anatomy of Marmoset Brain	172

1. Introduction

1.1. General remarks

Mice have made a great contribution to biomedical science and neuroscience because of its fertility and behavioral properties and development of breeding and genetic engineering techniques. There are, however, big differences in host responsiveness to pathogens, sensitivities to drugs, and characteristics of immune system, and therefore it is very difficult to interpret results from mice for clinical applications in humans. When we consider animal models for nervous and mental disease in particular, rodents are not ideal animals owing to huge differences in structure and functions of nervous system between rodents and humans. Thus, biomedical researchers have wanted primate models for nervous and mental disease that can express introduced genes. Rhesus monkeys (*Macaca mulatta*), the primate species most commonly used in biomedical science so far, have a far lower reproductive power than mice, and this low reproductive power prevented us from developing genetically modified primates.

Common marmosets (*Callithrix jacchus*) are small-bodied New World monkeys. They are endemic to a tropical forest in Brazil. The body weight of adult marmoset ranges 300 to 500 grams, and the body length is 20 to 25 cm. Unlike mice, the average life expectancy is 10 to 15 years and they are diurnal animals like humans. Marmosets have sensory abilities including visual abilities similar to humans. Recently, the small primates, common marmosets, have become the focus of attention of biomedical researchers because of its high reproductive power as well as its biosafety, ease of handling, and low cost for breeding.

Common marmosets reach sexual maturity at about 2 years. The ovarian cycle of common marmoset lasts about 28 days, and the gestation lasts about 5 months. Soon after parturition (within 10 days), female marmosets begin to cycle again and shortly thereafter become pregnant. The inter-birth interval is about 5 months and thus they give birth twice a year. Common marmosets usually give birth to twins; in captive, they can often have triplets. Therefore, female marmosets can have 40 to 60 offspring during their life. In contrast, female rhesus monkeys reach sexual maturity at about 5 years, usually have single birth, and usually have far fewer offspring (less than 10) during their life. This extremely high reproductive power can realize the development of genetically modified primate models for human disease, including nervous and mental disease.

In spite of the high potential of common marmosets as experimental animals for human disease, no brain atlas of marmoset using modern techniques has been available. One historical atlas was published but is now out of print (Stephan, Baron, Schwerdtfeger, 1980). Only one old paper published in a scientific journal is available (Saavedra, Mazzuchelli, 1968). Two brain atlas of marmoset were published in 2008; one (Palazzi, Bordier, 2008) includes slice images stained with acetyl choline esterase and those stained with cresyl violet and the other (Okano, Nomura, 2008) presented MR images acquired with a 7.0-T MR scanner with slice images stained with cresyl violet. We have realized the importance of generating a new brain atlas showing neurochemical characteristics in various regions for biomedical researches on the marmoset brain. We generated a brain atlas presenting serial sections immunostained with anti-calbindin, anti-calretinin, and anti-tyrosine hydroxylase antibodies as well as those stained with cresy violet. We also added MR images of marmoset brain to meet various demand for biomedical researches.

References

Saavedra J.P., Mazzuchelli A.L. (1968) A stereotaxic atlas of the brain of the marmoset (Hapale jacchus). Journal für Hirnforschung 11: 105-122.

Stephan H., Baron G., Schwerdtfeger W.K. (1980) The brain of the common marmoset (*Callithrix jacchus*): a stereotaxic atlas. Springer-Verlag, Berlin.

Palazzi X., Bordier N. (2008) The marmoset brain in stereotaxic coordinates. Springer, New York.

Okano H., Nomura T. (eds.) (2008) Brain MR atlas of common marmoset. Yodosha, Tokyo.

1.2. General construction of this atlas

Contents of Vol. 1

Coronal or parasagittal plates of serial sections stained with cresyl violet (Nissl staining) and the accompanying diagrams with labeling of structures. Corresponding plates of adjacent immunohistochemically-stained sections and MRI data in Vol. 2 are shown in the Tables of plates. Stereotaxic coordinates are indicated in the diagrams.

Contents of Vol. 2

Coronal or parasagittal plates of serial sections immunostained with anti-calbindin, anti-calretinin or anti-tyrosine hydroxylase antibody, and MRI data in the serial coronal or parasagittal sections. Corresponding plates of coronal or parasagittal sections stained with cresyl violet (Nissl staining) and the accompanying diagrams with labeling of structures in Vol. 1 are shown in the Tables of plates.

2. Materials and Methods

2.1. Histological procedures

All animal experiments were conducted in accordance with the NIH guidelines for the care and use of laboratory animals, and with the guidelines approved by the ethics committee for primate research of the National Center of Neurology and Psychiatry, Japan. An adult male marmoset, 1.6 years old and weighing 260 g was used for the preparation of coronal sections of the brain. Another adult male marmoset, 1.9 years old and weighing 320 g was used for the preparation of parasagittal sections of the brain. The marmosets were anesthetized with ketamine hydrochloride (30 mg/kg body weight, i.m.) and then a lethal dose of sodium pentobarbital (50 mg/kg, i.p.). Under deep anesthesia, the animals were perfused through the ascending aorta with 4% paraformaldehyde dissolved in 0.1 M phosphate-buffered saline (PBS, pH 7.4), by using a peristaltic pump. The brain was removed, and then postfixed at 4°C for 3 days with the same fixative. The fixed brains were embedded in 3% agar in PBS, and then sliced coronally into 100 µm sections with a microslicer (DTK-3000, DOSAKA EM, Kyoto, Japan) or sliced parasagittally into 150 µm sections with a microslicer (DTK-3000W, DOSAKA EM, Kyoto, Japan).

Nissl staining was performed on free-floating sections; sections were incubated in a solution of NeuroTrace 500/525 (fluorescent Nissl stain; Invitrogen, Carlsbad, CA, USA), 1:300 dilution in PBS containing 2% Triton X-100, for 2 hr at the room temperature. After rinse with PBS, the stained sections were mounted and cover-slipped on micro slide glass (76 x 52 mm, 0.8~1.0 mm thickness, Matsunamiglass, Osaka, Japan) with Fluoromount-G (SouthernBiotech, Birmingham, AL, USA).

Immunohistochemical staining was also performed on free-floating sections; the sections were pre-incubated with 10% normal goat serum at 4°C for 1 h. Then, sections were incubated with each primary antibody in PBS containing 2% Triton X-100 at 4°C overnight. The following antibodies were used in this study; anti-calbindin D-28k antibody (rabbit IgG, 1:2000 dilution; Cat. No. CB38a, Swant, Bellinzona, Switzerland), anti-calretinin antibody (rabbit IgG, 1:2000 dilution; Cat. No. 7699/4, Swant, Bellinzona, Switzerland), or monoclonal anti-tyrosine hydroxylase antibody (clone TH-16, mouse IgG, 1:1000 dilution; Cat. No. T2928, Sigma-Aldrich, St. Louis, MO, USA). Sectiones were then rinsed 8 times with PBS, and incubated with secondary antibody in PBS at 4°C for 5 h. Alexa Fluor 594 goat anti-rabbit IgG (1:500 dilution; Cat. No. A11012, Molecular Probes, Eugene, OR, USA) was used for anti-calbindin D-28k antibody or anti-calretinin antibody. Alexa Fluor 594 goat anti-mouse IgG (1:500 dilution; Cat. No. A11005, Molecular Probes, Eugene, OR, USA) was used for anti- tyrosine hydroxylase antibody. After incubation, the sections were rinsed 5 times with PBS. The stained sections were mounted on micro slide glass (76 x 52 mm, 0.8~1.0 mm thickness, Matsunami-glass, Osaka, Japan) and cover-slipped with Fluoromount-G (SouthernBiotech, Birmingham, AL, USA).

The above sections were examined with HS All-in-one fluorescence microscope BZ-9000 (Keyence Corporation, Osaka, Japan) in order to obtain the integrated images of wide area of the sections with high resolution following the manufacturer's instructions.

2.2. **MRI**

MRI data were acquired using a 3.0-T MRI scanner (Magnetom Trio; Siemens, Munich, Germany). A custom-built 4ch coil (Takashima Factory Ltd., Hino, Japan) was used for MR signal receiving. During MR imaging, marmosets were anesthetized with ketamine hydrochloride (7.5 mg/kg, i.m.) and pentobarbital sodium (15 mg/kg, i.p.), followed by an intramuscular injection of a mixture of ketamine hydrochloride (7.5 mg/kg) and medetomidine hydrochloride (0.15 mg/kg). The marmoset was fixed to a stereotaxic apparatus and set to the MR scanner. After manual shimming was conducted, high-resolu-

tion T1-weighted images (T1WIs) were obtained using the MPRAGE sequence. For coronal acquisition, parameters were set as TR = 2300 ms, TE = 2.38 ms, TI = 1000 ms, flip angle = 12°, matrix size = 192 \times 144, FoV = 66 \times 50 mm, slice thickness = 1 mm and total number of slices = 48. For sagittal acquisition, parameters were set as TR = 3000 ms, TE = 2.74 ms, TI = 1000 ms, flip angle = 12°, matrix size = 192 \times 192, FoV = 66 \times 66 mm, slice thickness = 1 mm and total number of slices = 48. The MR data were interpolated and voxel size was set to $0.174 \times 0.174 \times 1$ mm. In both coronal and sagittal acquisition, numbers of excitations were set at 8 and all MR volumes were averaged in order to increase signal to noise ratio. Each MR volume of coronal and sagittal acquisition was rotated as y-axis on the anterior and posterior commissure (AC-PC) line and z-axis on the interhemispheric fissure. Origins of MR volumes were set at the interaural line on the interhemispheric fissure. On coronal MRI figures, positive y values mean brain slices anterior to the interaural line and negative y values mean posterior slices, and right side of images is right side of brain in the neurological convention. On sagittal MRI figures, positive x values mean brain slices to the right of the interhemispheric fissure.

2.3. Stereotaxic coordinates

Coronal plane: the numbers at the bottom right of each drawing show the anteroposterior distance of the corresponding plate from the vertical plane passing through the interaural line and from the vertical plane passing through the bregma. The numbers on the left margin (on the ordinate) show the dorsoventral distance from the horizontal plane passing through the interaural line. The numbers on the bottom margins (on the abscissa) show the distance of structures from the midline (from the interhemispheric fissure).

Parasagittal plane: the number at the bottom right of each drawing shows the distance of the corresponding plate from the midline (from the interhemispheric fissure). The numbers on the left margin (on the ordinate) show the dorsoventral distance from the horizontal plane passing through the interaural line. The numbers on the bottom margin (on the abscissa) show the anteroposterior distance from the coronal plane passing through the interaural line.

3. Index of Abbreviations

2n	optic nerve	BLI	basolateral amygdaloid nucleus, intermedi-
3n	oculomotor nerve or its root		ate part
3N	oculomotor nucleus	BLVM	basolateral amygdaloid nucleus, ventrome-
3ND	oculomotor nucleus, dorsal part		dial part
3NV	oculomotor nucleus, ventral part	BM	basomedial amygdaloid nucleus
3V	3rd ventricle	BMMC	basomedial amygdaloid nucleus, magnocel-
4n	trochlear nerve or its root		lular part
4N	trochlear nucleus	BMPC	basomedial amygdaloid nucleus, parvicellu-
4V	4th ventricle		lar part
4VLR	lateral recess of the 4th ventricle	BMPCV	basomedial amygdaloid nucleus, parvicellu-
4x	trochlear decussation		lar part, ventral division
5N	trigeminal nucleus	bp	brachium pontis
6N	abducens nucleus	bsc	brachium of the superior colliculus
6n	root of abducens nerve	BST	bed nucleus of the stria terminalis
7N	facial nucleus	BSTIA	bed nucleus of the stria terminalis, intra-
7n	facial nerve		amygdaloid division
8n	vestibulocochlear nerve	CA	Cornu Ammonis in the hippocampus
10N	dorsal motor nucleus of vagus nerve	CA1	field CA1 of hippocampus
10n	vagus nerve	CA3	field CA3 of hippocampus
11N	accessory nucleus	CA4	field CA4 of hippocampus
12N	hypoglossal nucleus	cal	calcarine sulcus
12n	root of hypoglossal nerve	Cb1	cerebellar lobule 1
A1	primary auditory area	Cb2	cerebellar lobule 2
A2	second auditory area	Cb3	cerebellar lobule 3
AA	anterior amygdaloid area	Cb4	cerebellar lobule 4
AB	accessory basal nucleus	Cb5	cerebellar lobule 5
ac	anterior commissure	Cb6	cerebellar lobule 6
aca	anterior commissure, anterior part	Cb7	cerebellar lobule 7
Acb	accumbens nucleus	Cb8	cerebellar lobule 8
AcbSh	accumbens nucleus, shell	Cb9	cerebellar lobule 9
AcbC	accumbens nucleus, core	Cb10	cerebellar lobule 10
ACg	anterior cingulate gyrus	cc	corpus callosum
acp	anterior commissure, posterior part	Cd	caudate nucleus
AD	anterodorsal nucleus	Ce	central amygdaloid nucleus
ADH	dorsal hypothalamic area	CeL	central amygdaloid nucleus, lateral division
AH	Ammon's horn	CeM	central amygdaloid nucleus, medial
AHA	anterior hypothalamic area		division
AHi	amygdalohippocampal area	Cg	cingulate cortex
ALH	lateral hypothalamic area	CĞ	central gray
alv	alveus of the hippocampus	chp	choroid plexus
AM	anteromedial thalamic nucleus	CIC	central nucleus of the inferior colliculus
Amb	ambiguus nucleus	cic	commissure of the inferior colliculus
ans	ansiform lobule of the cerebellum	CIn	insularis cortex
AO	anterior olfactory nucleus	Cl	claustrum
APul	anterior pulvinar	CL	central lateral thalamic nucleus
Aq	aqueduct	CLi	caudal linear nucleus of the raphe
Arc	arcuate hypothalamic nucleus	CM	central medial thalamic nucleus
В	basal nucleus (Meynert)	CnF	cuneiform nucleus
BIC	nucleus of the brachium of the inferior	Co	cortical amygdaloid nucleus
	colliculus	Cop	copula of the pyramis
bic	brachium of the inferior colliculus	ср	cerebral peduncle, basal part
BL	basolateral amygdaloid nucleus	cr	corona radiata
BLD	basolateral amygdaloid nucleus, dorsal part	CR	central reticular nucleus
BLDL	basolateral amygdaloid nucleus, dorsolat-	Crus1	crus 1 of the ansiform lobule
	eral part	Crus2	crus 2 of the ansiform lobule

csc	commissure of the superior colliculus		nal band
CSG	cortex subgenualis	hf	hippocampal fissure
ctg	central tegmental tract	HR	hippocampus supracommissuralis
Cu	cuneate nucleus	ic	internal capsule
cu	cuneate fasciculus	IC	inferior colliculus
D3V	dorsal 3rd ventricle	icp	inferior cerebellar peduncle (restiform
DA	dorsoanterior extrastriate area		body)
dbc	decussatio brachii conjunctivi	IG	indusium griseum
DCIC	dorsal cortex of the inferior colliculus	IGP	internal globus pallidus
DEn	dorsal endopiriform nucleus	IH	infundibular nucleus
DG	dentate gyrus	ILL	intermediate nucleus of the lateral
DI	dorsointermediate visual area		lemniscus
dlf	dorsal longitudinal fasciculus	InfS	infundibular stem
DLG	dorsal lateral geniculate nucleus	InG	intermediate gray layer
DLL	dorsal nucleus of the lateral lemniscus	Int	interposed cerebellar nucleus
DM	dorsomedial hypothalamic nucleus	IntA	interposed cerebellar nucleus, anterior part
DM (visua		IntP	interposed cerebellar nucleus, posterior part
	dorsomedial visual area	IO	inferior olive
DM10	nucleus dorsalis motorius nervi vagi	IP	interpeduncular nucleus
DMC	dorsomedial hypothalamic nucleus compact	IPF	interpeduncular fossa
	part	IPul	inferior pulvinar
DPF1	dorsal paraflocculus	Irt	intermediate reticular nucleus
DpG	deep gray layer	IsC	Island of Calleja
DpMe	deep mesencephalic nucleus	ITG	inferotemporal gyrus
DR	dorsal raphe nucleus	La	lateral amygdaloid nucleus
dsc	dorsal spinocerebellar tract	Lat	lateral (dentate) cerebellar nucleus
DTg	dorsal tegmental nucleus	LC	locus coeruleus
ec	external capsule	lcsp	lateral corticospinal tract
ECIC	external cortex of the inferior colliculus	LD	lateral dorsal thalamic nucleus
ECu	external cuneate nucleus	LDSF	lateral dorsal thalamic nucleus, superficial
EGP	external globus pallidus		part
eml	external medullary lamina	lenf	lenticular fasciculus
EPL	external plexiform layer	lf	lateral fissure
Er	entorhinal cortex	lfp	longitudinal fasciculus of the pons
EW	Edinger-Westphal nucleus	LG	lateral geniculate nucleus
ex	extreme capsule	LH	lateral hypothalamic area
f	fornix	LHb	lateral habenular nucleus
FD	nucleus fasciculus diagonalis	11	lateral lemniscus
fi	fimbria of the hippocampus	LM	lateral mammillary nucleus
F1	flocculus	lml	lateral medullary lamina
flm	fasciculus longitudinalis medialis	LMol	lacunosum moleculare layer of the
flp	fasciculus longitudinalis pontis		hippocampus
Fr	frontal cortex	lo	lateral olfactory tract
fr	fasciculus retroflexus	LP	lateral posterior nucleus
FRP	formatio reticularis pontis	LPA	lateral preoptic area
FRT	formatio reticularis tegmenti	LPB	lateral parabrachial nucleus
FRTM	formatio reticularis tegmenti mesencephali	LPul	lateral pulvinar
g7	genu of the facial nerve	LRt	lateral reticular nucleus
GC	substantia grisea centralis	LS	lateral septal nucleus
Gi	gigantocellular reticular nucleus	LSD	lateral septal nucleus, dorsal part
Gl	glomerular layer	LSI	lateral septal nucleus, intermediate part
GL	granular cell layer	LSV	lateral septal nucleus, ventral part
GP	globus pallidus	Lu	stratum lucidum of the hippocampus
Gr	gracile nucleus	LV	lateral ventricle
gr	gracile fasciculus	LVeN	lateral vestibular nucleus
GrDG	granular layer of the dentate gyrus	lvsp	lateral vestibulospinal tract
Н	campus Foreli	M	mamillary nucleus
Hb	habenular nucleus	M1	primary motor area
HDB	nucleus of the horizontal limb of the diago-	m5	motor root of the trigeminal nerve

MC5	magnagallular layar of the could aring	Don	
MC3	magnocellular layer of the caudal spinal	Per	periamygdaloid nucleus
	trigeminal nucleus	PF	parafascicular thalamic nucleus
mcp	middle cerebellar peduncle	PFC	prefrontal cortex
MD	mediodorsal thalamic nucleus	PH	posterior hypothalamic area
MDC	mediodorsal thalamic nucleus, central part	Pi	pineal gland
MDD	mediodorsal thalamic nucleus, dorsal part	PM	paramedian lobule
MDL	mediodorsal thalamic nucleus, lateral part	PMn	paramedian reticular nucleus
MDM	mediodorsal thalamic nucleus, medial part	PMnR	paramedian raphe nucleus
MDR	medullary reticular nucleus	Pn	pontine nuclei
Me	medial amygdaloid nucleus	PnO	pontine reticular nucleus, oral part
ME	median eminence	POA	preoptic area
Me5N	mesencephalic trigeminal nucleus	POm	parietooccipital medial area
Med	medial (fastigial) cerebellar nucleus	PON	preoptic nucleus
mfb	medial forebrain bundle	Pp	peripeduncular nucleus
MG	medial geniculate nucleus	PP	posterior parietal cortex
MGD	medial geniculate nucleus, dorsal part	PPF1	posterior paraflocculus
MGM	medial geniculate nucleus, medial part	PPTgC	pedunculopontine tegmental nucleus, com-
MGV	medial geniculate nucleus, ventral part		pact part
MHb	medial habenular nucleus	PPTgD	pedunculopontine tegmental nucleus, dif-
Mi	mitral cell layer	8	fuse part
ML	medial mammillary nucleus, lateral part	PPY	prepiriform cortex
ml	medial lemniscus	Pr	prepositus nucleus
mlf	medial longitudinal fasciculus	PrCO	precentral orbital area
MM	medial mammillary nucleus, medial part	ProS	prosubiculum
mml	medial medullary lamina	PrS	presubiculum
MnR	median raphe nucleus	Pu	
Mo5N	motor trigeminal nucleus	ru Pul	putamen
Mol	molecular layer of the dentate gyrus	PUI	pulvinar nuclei
MPA			paraventricular thalamic nucleus
MPB	medial preoptic area	Pv	parietal ventral area
	medial parabrachial nucleus	py	pyramidal tract
MPr	motor and premotor cortex	pyrs	pyramis
MPul	medial pulvinar	pyx	pyramidal decussation
MS	medial septal nucleus	R	raphe nucleus
MT	middle temporal visual area	RA	radiatio acustica
MTc	middle temporal crescent visual area	rcc	rostrum of the corpus callosum
mtt	mamillothalamic tract	Re	reuniens thalamic nucleus
MVeN	medial vestibular nucleus	ReIC	recess of the inferior colliculus
	medial vestibular nucleus, magnocellular	RMC	red nucleus, magnocellular part
MVeNPC	medial vestibular nucleus, parvicellular	RMg	nucleus raphe magnus
NCS	central superior tegmental nucleus	RN	red nucleus
oc	olivocerebellar tract	RO	radiatio optica
OcC	occipital cortex	ROb	raphe obscurus nucleus
On	olivary nuclei	RP	nucleus raphe pallidus
ON	olfactory nerve layer	RPC	red nucleus, parvicellular part
opt	optic tract	RS	retrosplenial cortex
OT	olfactory tubercle	Rt	reticular thalamic nucleus
ots	occipitotemporal sulcus	RTP	nucleus reticularis tegmenti pontis
OV	olfactory ventricle	RtTg	reticulotegmental nucleus of the pons
OX	optic chiasm	S	subiculum
PaC	parietal cortex	S1	primary somatosensory area
PAG	periaqueductal gray	s5	sensory root of the trigeminal nerve
PaP	paraventricular hypothalamic nucleus,	SC	superior colliculus
	parvicellular part	SCG	superficial gray layer of superior colliculus
PaS	parasubiculum	SCN	suprachiasmatic nucleus
Pav	paraventricular hypothalamic nucleus	SCO	subcommissural organ
PBG	parabigeminal nucleus	scp	superior cerebellar peduncle (brachium
рс	posterior commissure	зер	conjunctivum)
PCRt	parvicellular reticular nucleus	Sen	
PCTg	paracollicular tegmentum	Sep SFi	septum septofimbrial nucleus
1015	paracomediar regimentum	SIT	septoninorial nucleus

SFO	subfornical organ		amygdala
SGe	supragenual nucleus	VAL	ventral anterior thalamic nucleus, lateral
SHi	septohippocampal nucleus		part
SI	substantia innominata	VAM	ventral anterior thalamic nucleus, medial
Sim	simple lobule		part
sm	stria medullaris of the thalamus	VC	ventral cochlear nucleus
SMH	supramamillary hypothalamic nucleus	VCA	ventral cochlear nucleus, anterior part
SMV	superior medullary velum	VCo	ventral cortical amygdaloid nucleus
SN	substantia nigra	VCP	ventral cochlear nucleus, posterior part
SNc	substantia nigra, compact	VDB	nucleus of the vertical limb of the diagonal
SNr	substantia nigra, reticular part		band
SO	superior olive	VeN	vestibular nuclei
Sol	solitary nucleus	VH	ventral horn
SON	supraoptic nucleus	vhc	ventral hippocampal commissure
SOR	supraoptic nucleus, retrochiasmatic part	VL	ventral lateral thalamic nucleus
SOX	supraoptic decussation	VLA	ventrolateral anterior extrastriate area
Sp5	spinal trigeminal nucleus	VLG	ventral lateral geniculate nucleus
sp5	spinal trigeminal tract	VLL	ventral lateral thalamic nucleus, lateral part
spth	spinothalamic tract	VLM	ventral lateral thalamic nucleus, medial part
st	stria terminalis	VLP	ventrolateral posterior extrastriate area
STG	superior temporal gyrus	VMH	ventromedial hypothalamic nucleus
STh	subthalamic nucleus	VP	ventral pallidum
STS	superior temporal sulcus	VPFl	ventral paraflocculus
SuM	supramammillary nucleus	VPL	ventral posterolateral thalamic nucleus
SVeN	superior vestibular nucleus	VPM	ventral posteromedial thalamic nucleus
TE	temporal cortex	VR	ventral raphe nucleus
TP	temporal polar cortex	vsc	ventral spinocerebellar tract
ts	tectospinal tract	VTA	ventral tegmental area
TuO	tuberculum olfactorium	xml	decussation of medial lemniscus
V1	primary visual cortex	xscp	decussation of the superior cerebellar
V2	second visual area		peduncle
VA	ventral anterior thalamic nucleus	ZI	zona incerta
VACo	ventral anterior cortical nucleus of the		

4. Tables of plates

coronal plane

tracing	Nissl staining	calbindin	calretinin	tyrosine hydroxylase	MRI
cTr-01	cNissl-01				
cTr-02	cNissl-02				MRI cor-0
cTr-03	cNissl-03				
cTr-04	cNissl-04				MRI cor-02
cTr-05	cNissl-05				MRI cor-03
cTr-06	cNissl-06	cCalb-06	cCR-06	cTH-06	MRI cor-04
cTr-07	cNissl-07				
cTr-08	cNissl-08	cCalb-08	cCR-08	cTH-08	
cTr-09	cNissl-09	cCalb-09	cCR-09	cTH-09	MRI cor-0:
cTr-10	cNissl-10				
cTr-11	cNissl-11	cCalb-11	cCR-11	cTH-11	
cTr-12	cNissl-12	cCalb-12	cCR-12	cTH-12	MRI cor-0
cTr-13	cNissl-13				
cTr-14	cNissl-14	cCalb-14	cCR-14	cTH-14	
cTr-15	cNissl-15	cCalb-15	cCR-15	cTH-15	MRI cor-0
cTr-16	cNissl-16				MRI cor-0
cTr-17	cNissl-17	cCalb-17	cCR-17	cTH-17	
cTr-18	cNissl-18	cCalb-18	cCR-18	cTH-18	MRI cor-09
cTr-19	cNissl-19	cCalb-19	cCR-19		
cTr-20	(cCalb-20)	cCalb-20	cCR-20		MRI cor-1
cTr-21	cNissl-21	cCalb-21	cCR-21	cTH-21	
cTr-22	cNissl-22	cCalb-22	cCR-22	cTH-22	
cTr-23	cNissl-23				
cTr-24	cNissl-24				MRI cor-1
cTr-25	cNissl-25				
cTr-26	cNissl-26	cCalb-26	cCR-26	cTH-26	
cTr-27	(cCalb-27)	cCalb-27		cTH-27	
cTr-28	(cCalb-28)	cCalb-28	cCR-28	cTH-28	
cTr-29	(cCalb-29)	cCalb-29	cCR-29		MRI cor-12
cTr-30	cNissl-30		cCR-30	cTH-30	
cTr-31	cNissl-31		cCR-31	cTH-31	
cTr-32	(cCalb-32)	cCalb-32	cCR-32	cTH-32	
cTr-33	cNissl-33	cCalb-33	cCR-33	cTH-33	MRI cor-13
cTr-34	cNissl-34	cCalb-34	cCR-34	cTH-34	
cTr-35	cNissl-35	cCalb-35	cCR-35	cTH-35	
cTr-36	cNissl-36	cCalb-36	cCR-36	cTH-36	MRI cor-14
cTr-37	cNissl-37	cCalb-37	cCR-37	cTH-37	
cTr-38	cNissl-38	cCalb-38	cCR-38	cTH-38	
cTr-39	cNissl-39	cCalb-39			MRI cor-1:
cTr-40	cNissl-40	cCalb-40		cTH-40	
cTr-41	cNissl-41			cTH-41	
cTr-42	cNissl-42				


coronal plane

tracing	Nissl staining	calbindin	calretinin	tyrosine hydroxylase	MRI
cTr-43	cNissl-43	cCalb-43	cCR-43	cTH-43	MRI cor-16
cTr-44	cNissl-44	cCalb-44	cCR-44	cTH-44	
cTr-45	cNissl-45	cCalb-45	cCR-45	cTH-45	
cTr-46	cNissl-46	cCalb-46	cCR-46	cTH-46	MRI cor-17
cTr-47	cNissl-47	cCalb-47	cCR-47	cTH-47	
cTr-48	cNissl-48	cCalb-48	cCR-48	cTH-48	
cTr-49	(cCalb-49)	cCalb-49	cCR-49	cTH-49	MRI cor-18
cTr-50	cNissl-50	cCalb-50	cCR-50		
cTr-51	cNissl-51	cCalb-51	cCR-51		
cTr-52	cNissl-52	cCalb-52	cCR-52		MRI cor-19
cTr-53	cNissl-53	cCalb-53	cCR-53		MRI cor-20
cTr-54	cNissl-54		cCR-54		MRI cor-21
cTr-55	cNissl-55		cCR-55		
cTr-56	cNissl-56	cCalb-56	cCR-56		MRI cor-22
cTr-57	cNissl-57	cCalb-57	cCR-57		
cTr-58	cNissl-58	cCalb-58	cCR-58		MRI cor-23
cTr-59	cNissl-59	cCalb-59	cCR-59		
cTr-60	cNissl-60	cCalb-60	cCR-60		
cTr-61	cNissl-61	cCalb-61	cCR-61		MRI cor-24
cTr-62	cNissl-62	cCalb-62	cCR-62		
cTr-63	cNissl-63	cCalb-63	cCR-63		MRI cor-25
cTr-64	cNissl-64				MRI cor-26
cTr-65	cNissl-65	cCalb-65	cCR-65		MRI cor-23
cTr-66	cNissl-66	cCalb-66	cCR-66		MRI cor-26

parasagittal plane

tracing	Nissl staining	calbindin	calretinin	tyrosine hydroxylase	MRI
psTr-01	psNissl-01	psCalb-01	psCR-01		MRI psg-01
psTr-02	psNissl-02	psCalb-02	psCR-02	psTH-02	MRI psg-02
psTr-03	psNissl-03	psCalb-03	psCR-03	psTH-03	MRI psg-03
psTr-04	psNissl-04	psCalb-04		psTH-04	
psTr-05	psNissl-05	psCalb-05	psCR-05	psTH-05	MRI psg-04
psTr-06	psNissl-06	psCalb-06	psCR-06	psTH-06	
psTr-07	psNissl-07	psCalb-07	psCR-07	psTH-07	MRI psg-05
psTr-08	psNissl-08	psCalb-08	psCR-08	psTH-08	MRI psg-06
psTr-09	psNissl-09	psCalb-09	psCR-09		MRI psg-07
psTr-10	psNissl-10	psCalb-10	psCR-10	psTH-10	
psTr-11	psNissl-11	psCalb-11	psCR-11	psTH-11	MRI psg-08
psTr-12	psNissl-12	psCalb-12	psCR-12	psTH-12	MRI psg-09
psTr-13	psNissl-13	psCalb-13	psCR-13		MRI psg-10~11
psTr-14	psNissl-14	psCalb-14			MRI psg-12


5.	Plates:	Nissl-stained	sections	and	corresponding	tracing
----	----------------	---------------	----------	-----	---------------	---------



EPL, external plexiform layer; GI, glomerular layer; GL, granular cell layer; Mi, mitral cell layer; ON, olfactory nerve layer; OV, olfactory ventricle.


cNissl-01






EPL, external plexiform layer; GI, glomerular layer; GL, granular cell layer; Mi, mitral cell layer; ON, olfactory nerve layer; OV, olfactory ventricle.


cNissI-02






EPL, external plexiform layer; GI, glomerular layer; GL, granular cell layer; Mi, mitral cell layer; ON, olfactory nerve layer; OV, olfactory ventricle.

cNissI-03





8Ad, 8Av, 8B, 12/45, 13, 32, 46, Brodmann's parcellation of the cortex; Cl, claustrum; lo, lateral olfactory tract; Prefr, prefrontal cortex.

Brodmann's parcellation of the marmoset cortex is according to the following literature.

Burman KJ, Palmer SM, Gamberini M, Rosa MGP. 2006. Cytoarchitectonic subdivisions of the dorsolateral frontal cortex of the marmoset monkey (*Callithrix jacchus*), and their projections to dorsal visual areas. J Comp Neurol 495: 149-172.

Burman KJ, Lui LL, Rosa MGP, Bourne JA. 2007. Development of non-phosphorylated neurofilament protein expression in neurons of the New World monkey dorsolateral frontal cortex. Eur J Neurosci 25: 1767-1779. Burman KJ, Palmer SM, Gamberini M, Spitzer MW, Rosa MGP. 2008. Anatomical and physiological definition of the motor cortex of the marmoset monkey. J Comp Neurol 506: 860-876.


Burish MJ, Stepniewska I, Kaas JH. 2008. Microstimulation and architectonics of frontoparietal cortex in common marmosets (*Callithrix jacchus*). J Comp Neurol 507: 1151-1168.


Roberts AC, Tomic DL, Parkinson CH, Roeling TA, Cutter DJ, Robbins TW, Everitt BJ. 2007.

Forebrain connectivity of the prefrontal cortex in the marmoset monkey (*Callithrix jacchus*): an anterograde and retrograde tract-tracing study. J Comp Neurol 502: 86-112.

Rosa MGP, Tweedale R. 2000. Visual areas in lateral and ventral extrastriate cortices of the marmoset monkey. J Comp Neurol 422: 621-651.


cNissl-04






8Ad, 8Av, 8B, 12/45, 13, 24, 25, 32, 46, Brodmann's parcellation of the cortex; CI, claustrum; lo, lateral olfactory tract; Prefr, prefrontal cortex.


cNissI-05






6dr, 6m, 8Av, 12/45, 13, 24, 25, 32, Brodmann's parcellation of the cortex; Cl, claustrum; Fr, frontal cortex; lo, lateral olfactory tract.

cNissl-06






6dr, 6m, 8Av, 12/45, 13, 24, 25, Brodmann's parcellation of the cortex; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; Cln, insularis cortex; Cl, claustrum; Fr, frontal cortex; lo, lateral olfactory tract; LV, lateral ventricle; PrCO, precentral orbital area.


cNissl-07






6dr, 6m, 8Av, 12/45, 13, 24, 25, Brodmann's parcellation of the cortex; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; Cln, insularis cortex; Fr, frontal cortex; lo, lateral olfactory tract; LV, lateral ventricle; PrCO, precentral orbital area.


cNissl-08






6dr, 6m, 8Av, 12/45, 13, 24, 25, Brodmann's parcellation of the cortex; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; Cln, insularis cortex; Fr, frontal cortex; LV, lateral ventricle.


cNissl-09





6dr, 6m, 8Av, 12/45, 13, 24, 25, Brodmann's parcellation of the cortex; Acb, accumbens nucleus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; Cln, insularis cortex; Cl, claustrum; CSG, cortex subgenualis; ec, external capsule; Fr, frontal cortex; ic, internal capsule; IsC, Island of Calleja; PPY, prepiriform cortex; Pu. Putamen.






3a, 6d, 6m, 6v, 12/45, 13, 24, Brodmann's parcellation of the cortex; Acb, accumbens nucleus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; Cln, insularis cortex; Cl, claustrum; cr, corona radiata; ec, external capsule; Fr, frontal cortex; HDB, nucleus of the horizontal limb of the diagonal band; ic, internal capsule; IsC, Island of Calleja; LSD, lateral septal nucleus, dorsal part; LV, lateral ventricle; MS, medial septal nucleus; OT, olfactory tubercle; Pu, putamen; VDB, nucleus of the vertical limb of the diagonal band.


cNissI-11







3a, 6d, 6m, 6v, 12/45, 24, Brodmann's parcellation of the cortex; Acb, accumbens nucleus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; Cln, insularis cortex; Cl, claustrum; ec, external capsule; Fr, frontal cortex; HDB, nucleus of the horizontal limb of the diagonal band; ic, internal capsule; IsC, Island of Calleja; LS, lateral septal nucleus; LV, lateral ventricle; MS, medial septal nucleus; Pu, putamen; VDB, nucleus of the vertical limb of the diagonal band.







3a, 3b, 6d, 6m, 6v, 12/45, 24, Brodmann's parcellation of the cortex; Acb, accumbens nucleus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; Cln, insularis cortex; Cl, claustrum; cr, corona radiata; ec, external capsule; Fr, frontal cortex; HDB, nucleus of the horizontal limb of the diagonal band; ic, internal capsule; IsC, Island of Calleja; LS, lateral septal nucleus; LV, lateral ventricle; MS, medial septal nucleus; Pu, putamen; S1, primary somatosensory area; VDB, nucleus of the vertical limb of the diagonal band.







3a, 3b, 6d, 6m, 6v, 24, Brodmann's parcellation of the cortex; Acb, accumbens nucleus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; Cln, insularis cortex; Cl, claustrum; cr, corona radiata; ec, external capsule; Fr, frontal cortex; HDB, nucleus of the horizontal limb of the diagonal band; ic, internal capsule; IsC, Island of Calleja; If, lateral fissure; LS, lateral septal nucleus; LV, lateral ventricle; MS, medial septal nucleus; OT, olfactory tubercle; Pu, putamen; S1, primary somatosensory area; TP, temporal polar cortex; VDB, nucleus of the vertical limb of the diagonal band.






3a, 3b, 4, 6d, 6m, 6v, 13, 24, Brodmann's parcellation of the cortex; AA, anterior amygdaloid area; Acb, accumbens nucleus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; Cln, insularis cortex; Cl, claustrum; ec, external capsule; ex, extreme capsule; Fr, frontal cortex; HDB, nucleus of the horizontal limb of the diagonal band; ic, internal capsule; If, lateral fissure; LSD, lateral septal nucleus, dorsal part; LSI, lateral septal nucleus, intermediate part; LSV, lateral septal nucleus, ventral part; LV, lateral ventricle; M1, primary motor area; MS, medial septal nucleus; Pu, putamen; S1, primary somatosensory area; TP, temporal polar cortex; VDB, nucleus of the vertical limb of the diagonal band.






3a, 3b, 4, 6d, 6m, 6v, 13, 24, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; AA, anterior amygdaloid area; Acb, accumbens nucleus; BL, basolateral amygdaloid nucleus; BM, basomedial amygdaloid nucleus; BST, bed nucleus of the stria terminalis; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; Cln, insularis cortex; Co, cortical amygdaloid nucleus; Er, entorhinal cortex; Fr; frontal cortex; HDB, nucleus of the horizontal limb of the diagonal band; ic, internal capsule; If, lateral fissure; La, lateral amygdaloid nucleus; LPA, lateral preoptic area; LSD, lateral septal nucleus, dorsal part; LSI, lateral septal nucleus, intermediate part; LV, lateral ventricle; M1, primary motor area; Me, medial amygdaloid nucleus; MPA, medial preoptic area; MS, medial septal nucleus; ox, optic chiasm; Pu, putamen; S1, primary somatosensory area; SHi, septohippocampal nucleus; SON, supraoptic nucleus; TP, temporal polar cortex.






3a, 3b, 4, 13, 20, 22, 24, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; AA, anterior amygdaloid area; ac, anterior commissure; BL, basolateral amygdaloid nucleus; BM, basomedial amygdaloid nucleus; BST, bed nucleus of the stria terminalis; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; Cln, insularis cortex; Cl, claustrum; Er, entorhinal cortex; Fr; frontal cortex; GP, globus pallidus; ic, internal capsule; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; If, lateral fissure; LSD, lateral septal nucleus, dorsal part; LSI, lateral septal nucleus, intermediate part; LV, lateral ventricle; M1, primary motor area; Me, medial amygdaloid nucleus; MPA, medial preoptic area; MS, medial septal nucleus; ox, optic chiasm; PaC, parietal cortex; Pu, putamen; Pv, parietal ventral area; S1, primary somatosensory area; SCN, suprachiasmatic nucleus; SON, supraoptic nucleus; STG, superior temporal gyrus; TE, temporal cortex.




3a, 3b, 4, 13, 20, 22, 24, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; AB, accessory basal nucleus; ac, anterior commissure; acp, anterior commissure, posterior part; BL, basolateral amygdaloid nucleus; BM, basomedial amygdaloid nucleus; BST, bed nucleus of the stria terminalis; cc, corpus callosum; Cd, caudate nucleus; Ce, central amygdaloid nucleus; Cg, cingulate cortex; Cln, insularis cortex; Cl, claustrum; cr, corona radiata; Er, entorhinal cortex; f, fornix; FD, nucleus fasciculus diagonalis; Fr; frontal cortex; GP, globus pallidus; ic, internal capsule; IG, indusium griseum; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; If, lateral fissure; LPA, lateral preoptic area; LS, lateral septal nucleus; LV, lateral ventricle; M1, primary motor area; Me, medial amygdaloid nucleus; MPA, medial preoptic area; ox, optic chiasm; PaC, parietal cortex; Pu, putamen; Pv, parietal ventral area; S1, primary somatosensory area; SCN, suprachiasmatic nucleus; SON, supraoptic nucleus; st, stria terminalis; STG, superior temporal gyrus; TE, temporal cortex.






3a, 3b, 4, 13, 20, 22, 24, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; AB, accessory basal nucleus; ac, anterior commissure; AHA, anterior hypothalamic area; BL, basolateral amygdaloid nucleus; BM, basomedial amygdaloid nucleus; cc, corpus callosum; Cd, caudate nucleus; CeL, central amygdaloid nucleus, lateral division; CeM, central amygdaloid nucleus, medial division; Cg, cingulate cortex; Cln, insularis cortex; Co, cortical amygdaloid nucleus; Er, entorhinal cortex; f, fornix; Fr; frontal cortex; GP, globus pallidus; ic, internal capsule; IG, indusium griseum; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; If, lateral fissure; LS, lateral septal nucleus; LV, lateral ventricle; M1, primary motor area; Me, medial amygdaloid nucleus; MS, medial septal nucleus; ox, optic chiasm; PaC, parietal cortex; Pav, paraventricular hypothalamic nucleus; Pu, putamen; Pv, parietal ventral area; S1, primary somatosensory area; SCN, suprachiasmatic nucleus; SI, substantia innominata; SON, supraoptic nucleus; st, stria terminalis; STG, superior temporal gyrus; TE, temporal cortex.






3a, 3b, 4, 13, 20, 22, 24, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; acp, anterior commissure, posterior part; AHA, anterior hypothalamic area; BLD, basolateral amygdaloid nucleus, dorsal part; BLI, basolateral amygdaloid nucleus, intermediate part; BM, basomedial amygdaloid nucleus; BST, bed nucleus of the stria terminalis; cc, corpus callosum; Cd, caudate nucleus; CeL, central amygdaloid nucleus, lateral division; CeM, central amygdaloid nucleus, medial division; Cg, cingulate cortex; CIn, insularis cortex; CI, claustrum; cr, corona radiata; EGP, external globus pallidus; f, fornix; Fr; frontal cortex; ic, internal capsule; IGP, internal globus pallidus; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; If, lateral fissure; LV, lateral ventricle; M1, primary motor area; Me, medial amygdaloid nucleus; ox, optic chiasm; PaC, parietal cortex; Pav, paraventricular hypothalamic nucleus; Pu, putamen; Pv, parietal ventral area; S1, primary somatosensory area; SCN, suprachiasmatic nucleus; SON, supraoptic nucleus; STG, superior temporal gyrus; TE, temporal cortex; VA, ventral anterior thalamic nucleus; VP, ventral pallidum.

cCalb-20



3b, 4, 13, 20, 22, 24, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; acp, anterior commissure, posterior part; ADH, dorsal hypothalamic area; Arc, arcuate hypothalamic nucleus; B, basal nucleus (Meynert); BLD, basolateral amygdaloid nucleus,dorsal part; BLI, basolateral amygdaloid nucleus,intermediate part; BLVM, basolateral amygdaloid nucleus,ventromedial part; BM, basomedial amygdaloid nucleus; cc, corpus callosum; Cd, caudate nucleus; CeL, central amygdaloid nucleus,lateral division; CeM, central amygdaloid nucleus,medial division; Cg, cingulate cortex; CIn, insularis cortex; CI, claustrum; Co, cortical amygdaloid nucleus; cr, corona radiata; EGP, external globus pallidus; Er, entorhinal cortex; f, fornix; Fr; frontal cortex; ic, internal capsule; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; If, lateral fissure; ImI, lateral medullary lamina; LV, lateral ventricle; M1, primary motor area; ME, median eminence; Me, medial amygdaloid nucleus; opt, optic tract; PaC, parietal cortex; Pav, paraventricular hypothalamic nucleus; Per, periamygdaloid nucleus; Pu, putamen; S1, primary somatosensory area; SON, supraoptic nucleus; STG, superior temporal gyrus; TE, temporal cortex; VA, ventral anterior thalamic nucleus; VP, ventral pallidum.






1+3, 13, 20, 22, 24, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; acp, anterior commissure, posterior part; ADH, dorsal hypothalamic area; ALH, lateral hypothalamic area; Arc, arcuate hypothalamic nucleus; B, basal nucleus (Meynert); BL, basolateral amygdaloid nucleus; BM, basomedial amygdaloid nucleus; cc, corpus callosum; Cd, caudate nucleus; Ce, central amygdaloid nucleus; Cg, cingulate cortex; Cln, insularis cortex; Cl, claustrum; Co, cortical amygdaloid nucleus; ec, external capsule; Er, entorhinal cortex; ex, extreme capsule; f, fornix; GP, globus pallidus; ic, internal capsule; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; LV, lateral ventricle; Me, medial amygdaloid nucleus; opt, optic tract; Pav, paraventricular hypothalamic nucleus; Pac, parietal cortex, Per, periamygdaloid nucleus; Pu, putamen; STG, superior temporal gyrus; TE, temporal cortex; VA, ventral anterior thalamic nucleus; VMH, ventromedial hypothalamic nucleus.


cNissI-22







1+3, 13, 20, 22, 24, 41, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A1, primary auditory area; acp, anterior commissure, posterior part; Arc, arcuate hypothalamic nucleus; BLD, basolateral amygdaloid nucleus, dorsal part; BLI, basolateral amygdaloid nucleus, intermediate part; BLVM, basolateral amygdaloid nucleus, ventromedial part; BM, basomedial amygdaloid nucleus; BST, bed nucleus of the stria terminalis: cc, corpus callosum; Cd, caudate nucleus; CeL, central amygdaloid nucleus, lateral division; CeM, central amygdaloid nucleus, medial division; Cg, cingulate cortex; Cln, insularis cortex; Cl, claustrum; CM, central medial thalamic nucleus; Co, cortical amygdaloid nucleus; cr, corona radiata; DM, dorsomedial hypothalamic nucleus; EGP, external globus pallidus; Er, entorhinal cortex; f, fornix; ic, internal capsule; IG, indusium griseum; IGP, internal globus pallidus; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; If, lateral fissure; LH, lateral hypothalamic area; Iml, lateral medullary lamina; LV, lateral ventricle; Me, medial amygdaloid nucleus; mml, medial medullary lamina; opt, optic tract; PaC, parietal cortex; PH, posterior hypothalamic area; Pu, putamen; SHi, septohippocampal nucleus; SON, supraoptic nucleus; st, stria terminalis; STG, superior temporal gyrus; TE, temporal cortex; VMH, ventromedial hypothalamic nucleus.


cNissI-23






5,13, 20, 22, 23, 41, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A1, primary auditory area; acp, anterior commissure, posterior part; AH, Ammon's horn; BL, basolateral amygdaloid nucleus; BM, basomedial amygdaloid nucleus; cc, corpus callosum; Cd, caudate nucleus; CeL, central amygdaloid nucleus, lateral division; CeM, central amygdaloid nucleus, medial division; Cg, cingulate cortex; CIn, insularis cortex; Co, cortical amygdaloid nucleus; DG, dentate gyrus; EGP, external globus pallidus; Er, entorhinal cortex; f, fornix; ic, internal capsule; IGP, internal globus pallidus; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; LD, lateral dorsal thalamic nucleus; lenf, lenticular fasciculus; If, lateral fissure; LH, lateral hypothalamic area; LM, lateral mammillary nucleus; LV, lateral ventricle; MD, mediodorsal thalamic nucleus; Me, medial amygdaloid nucleus; MM, medial mammillary nucleus, medial part; opt, optic tract; PH, posterior hypothalamic area; PP, posterior parietal cortex; Pu, putamen; PV, paraventricular thalamic nucleus; Rt, reticular thalamic nucleus; S2, subiculum; SON, supraoptic nucleus; S2, secondary somatosensory area; STG, superior temporal gyrus; TE, temporal cortex; VAL, ventral anterior thalamic nucleus, lateral part; VAM, ventral anterior thalamic nucleus, medial part; VL, ventral lateral thalamic nucleus.






5,13, 20, 22, 23, 41, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A1, primary auditory area; acp, anterior commissure, posterior part; AH, Ammon's horn; BL, basolateral amygdaloid nucleus; BM, basomedial amygdaloid nucleus; cc, corpus callosum; Cd, caudate nucleus; Ce, central amygdaloid nucleus; Cg, cingulate cortex; CIn, insularis cortex; CL, central lateral thalamic nucleus; Co, cortical amygdaloid nucleus; DG, dentate gyrus; DM, dorsomedial hypothalamic nucleus; EGP, external globus pallidus; Er, entorhinal cortex; f, fornix; ic, internal capsule; IGP, internal globus pallidus; IH, infundibular nucleus; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; LD, lateral dorsal thalamic nucleus; LH, lateral hypothalamic area; LV, lateral ventricle; M, mamillary nucleus; MD, mediodorsal thalamic nucleus; Me, medial amygdaloid nucleus; opt, optic tract; PH, posterior hypothalamic area; PP, Posterior parietal cortex; Pu, putamen; Rt, reticular thalamic nucleus; S, subiculum; SON, supraoptic nucleus; S2, secondary somatosensory area; STG, superior temporal gyrus; TE, temporal cortex; VAL, ventral anterior thalamic nucleus, lateral part; VAM, ventral anterior thalamic nucleus, medial part; VL, ventral lateral thalamic nucleus.







5,13, 20, 22, 23, 41, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A1, primary auditory area; Arc, arcuate hypothalamic nucleus; B, basal nucleus (Meynert); BL, basolateral amygdaloid nucleus; BLD, basolateral amygdaloid nucleus; BLD, basolateral amygdaloid nucleus, CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Ce, central amygdaloid nucleus; Cg, cingulate cortex; Cln, insularis cortex; CL, central lateral thalamic nucleus; CM, central medial thalamic nucleus; Co, cortical amygdaloid nucleus; DG, dentate gyrus; DM, dorsomedial hypothalamic nucleus; EGP, external globus pallidus; Er, entorhinal cortex; f, fornix; ic, internal capsule; IGP, internal globus pallidus; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; LD, lateral dorsal thalamic nucleus; If, lateral fissure; LH, lateral hypothalamic area; LV, lateral ventricle; M, mamillary nucleus; MD, mediodorsal thalamic nucleus; Me, medial amygdaloid nucleus; opt, optic tract; PH, posterior hypothalamic area; PP, posterior parietal cortex; Pu, putamen; Rt, reticular thalamic nucleus; S, subiculum; S2, secondary somatosensory area; STG, superior temporal gyrus; TE. temporal cortex; VA, ventral anterior thalamic nucleus; VL, ventral lateral thalamic nucleus; ZI, zona incerta.






5,13, 20, 22, 23, 41, 42, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A1, primary auditory area; A2, secondary auditory region; acp, anterior commissure, posterior part; Arc, arcuate hypothalamic nucleus; BL, basolateral amygdaloid nucleus; BM, basomedial amygdaloid nucleus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Ce, central amygdaloid nucleus; Cg, cingulate cortex; Cln, insularis cortex; CL, central lateral thalamic nucleus; CM, central medial thalamic nucleus; Co, cortical amygdaloid nucleus; DG, dentate gyrus; DM, dorsomedial hypothalamic nucleus; EGP, external globus pallidus; Er, entorhinal cortex; f, fornix; ic, internal capsule; IGP, internal globus pallidus; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; LD, lateral dorsal thalamic nucleus; If, lateral fissure; LH, lateral hypothalamic area; LV, lateral ventricle; M, mamillary nucleus; MD, mediodorsal thalamic nucleus; Me, medial amygdaloid nucleus; opt, optic tract; PP, posterior parietal cortex; Pu, putamen; Rt, reticular thalamic nucleus; VL, ventral lateral thalamic nucleus; ZI, zona incerta.







5,13, 20, 22, 23, 41, 42, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A1, primary auditory area; A2, secondary auditory region; acp, anterior commissure, posterior part; Arc, arcuate hypothalamic nucleus; BL, basolateral amygdaloid nucleus; BM, basomedial amygdaloid nucleus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Ce, central amygdaloid nucleus; Cg, cingulate cortex; CIn, insularis cortex: CL, central lateral thalamic nucleus; CM, central medial thalamic nucleus; DG, dentate gyrus; DM, dorsomedial hypothalamic nucleus; EGP, external globus pallidus; Er, entorhinal cortex; f, fornix; ic, internal capsule; IGP, internal globus pallidus; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; LD, lateral dorsal thalamic nucleus; lenf, lenticular fasciculus; If, lateral fissure; LH, lateral hypothalamic area; LM, lateral mammillary nucleus; LV, lateral ventricle; MD, mediodorsal thalamic nucleus; Me, medial amygdaloid nucleus; MM, medial mammillary nucleus, medial part; opt, optic tract; PP, posterior parietal cortex; Pu, putamen; Rt, reticular thalamic nucleus; S, subiculum; sm, stria medullaris of the thalamus; SON, supraoptic nucleus; STG, superior temporal gyrus; TE, temporal cortex; VA, ventral anterior thalamic nucleus; VL, ventral lateral thalamic nucleus; ZI, zona incerta.







5,13, 20, 22, 23, 41, 42, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A1, primary auditory area; A2, secondary auditory region; acp, anterior commissure, posterior part; Arc, arcuate hypothalamic nucleus; BL, basolateral amygdaloid nucleus; BM, basomedial amygdaloid nucleus; CA, Cornu Ammonis; cc, corpus callosum; Cd, caudate nucleus; Ce, central amygdaloid nucleus; Cg, cingulate cortex; Cln, insularis cortex: CM, central medial thalamic nucleus; DG, dentate gyrus; DM, dorsomedial hypothalamic nucleus; EGP, external globus pallidus; Er, entorhinal cortex; f, fornix; ic, internal capsule; IGP, internal globus pallidus; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; LD, lateral dorsal thalamic nucleus; If, lateral fissure; LH, lateral hypothalamic area; LM, lateral mammillary nucleus; LV, lateral ventricle; MD, mediodorsal thalamic nucleus; Me, medial amygdaloid nucleus; MM, medial mammillary nucleus, medial part; opt, optic tract; PP, posterior parietal cortex; Pu, putamen; Rt, reticular thalamic nucleus; S, subiculum; SON, supraoptic nucleus; STG, superior temporal gyrus; TE, temporal cortex; VA, ventral anterior thalamic nucleus; VL, ventral lateral thalamic nucleus; ZI, zona incerta.







5,13, 20, 22, 23, 41, 42, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A1, primary auditory area; A2, secondary auditory region; acp, anterior commissure, posterior part; Arc, arcuate hypothalamic nucleus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; Cln, insularis cortex: Cl, claustrum; CL, central lateral thalamic nucleus; CM, central medial thalamic nucleus; DG, dentate gyrus; DM, dorsomedial hypothalamic nucleus; EGP, external globus pallidus; Er, entorhinal cortex; f, fornix; ic, internal capsule; IGP, internal globus pallidus; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; LD, lateral dorsal thalamic nucleus; lenf, lenticular fasciculus; If, lateral fissure; LH, lateral hypothalamic area; Iml, lateral medullary lamina; LV, lateral ventricle; M, mamillary nucleus; MD, mediodorsal thalamic nucleus; mml, medial medullary lamina; opt, optic tract; PH, posterior hypothalamic area; PP, Posterior parietal cortex; Pu, putamen; Rt, reticular thalamic nucleus; S, subiculum; SON, supraoptic nucleus; st, stria terminalis; STG, superior temporal gyrus; TE, temporal cortex; VL, ventral lateral thalamic nucleus; ZI, zona incerta.






5, 20, 22, 23, 42, 50, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A2, secondary auditory region; Arc, arcuate hypothalamic nucleus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; CL, central lateral thalamic nucleus; CM, central medial thalamic nucleus; cp, cerebral peduncle,basal part; DG, dentate gyrus; ec, external capsule; EGP, external globus pallidus; Er, entorhinal cortex; f, fornix; ic, internal capsule; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; LD, lateral dorsal thalamic nucleus; If, lateral fissure; LH, lateral hypothalamic area; Iml, lateral medulary lamina; LV, lateral ventricle; MD, mediodorsal thalamic nucleus; ML, medial mammillary nucleus,lateral part; MM, medial mammillary nucleus,medial part; mml, medial medullary lamina; opt, optic tract; PH, posterior hypothalamic area; PP, posterior parietal cortex; Pu, putamen; Rt, reticular thalamic nucleus; S, subiculum; st, stria terminalis; STG, superior temporal gyrus; TE, temporal cortex; VL, ventral lateral thalamic nucleus; VPL, ventral posterolateral thalamic nucleus; ZI, zona incerta.






5, 20, 21, 22, 23, 42, 50, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A2, secondary auditory region; acp, anterior commissure, posterior part; Arc, arcuate hypothalamic nucleus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; CL, central lateral thalamic nucleus; CM, central medial thalamic nucleus; cp, cerebral peduncle,basal part; DG, dentate gyrus; ec, external capsule; Er, entorhinal cortex; f, fornix; GP, globus pallidus; ic, internal capsule; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; LD, lateral dorsal thalamic nucleus; If, lateral fissure; LV, lateral ventricle; MD, mediodorsal thalamic nucleus; ML, medial mammillary nucleus,lateral part; MM, medial mammillary nucleus,medial part; mtt, mamillothalamic tract; opt, optic tract; PH, posterior hypothalamic area; PP, posterior parietal cortex; Pu, putamen; Rt, reticular thalamic nucleus; S, subiculum; st, stria terminalis; STG, superior temporal gyrus; STh, subthalamic nucleus; VPM, ventral posterolateral thalamic nucleus; VPM, ventral posteromedial thalamic nucleus; ZI, zona incerta.






5, 20, 21, 22, 23, 42, 50, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A2, secondary auditory region; ADH, dorsal hypothalamic area; ALH, lateral hypothalamic area; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; CL, central lateral thalamic nucleus; CM, central medial thalamic nucleus; cp, cerebral peduncle,basal part; DG, dentate gyrus; EGP, external globus pallidus; Er, entorhinal cortex; f, fornix; ic, internal capsule; IGP, internal globus pallidus; ITG, inferotemporal gyrus; La, lateral amygdaloid nucleus; LD, lateral dorsal thalamic nucleus; lenf, lenticular fasciculus; If, lateral fissure; LV, lateral ventricle; MD, mediodorsal thalamic nucleus; ML, medial mammillary nucleus,medial part; mtt, mamillothalamic tract; opt, optic tract; PF, parafascicular thalamic nucleus; PP, posterior parietal cortex; Pu, putamen; Rt, reticular thalamic nucleus; SuM, supramammillary nucleus; TE, temporal cortex; VL, ventral lateral thalamic nucleus; VPL, ventral posterolateral thalamic nucleus; VPL, ventral posterolateral thalamic nucleus; ZI, zona incerta.







5, 20, 21, 22, 23, 42, 50, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A2, secondary auditory region; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; CL, central lateral thalamic nucleus; Cl, claustrum; CM, central medial thalamic nucleus; cp, cerebral peduncle,basal part; DG, dentate gyrus; DM, dorsomedial hypothalamic nucleus; ec, external capsule; EGP, external globus pallidus; Er, entorhinal cortex; f, fornix; ic, internal capsule; IGP, internal globus pallidus; ITG, inferotemporal gyrus; LD, lateral dorsal thalamic nucleus; If, lateral fissure; LH, lateral hypothalamic area, lml, lateral medullary lamina; LV, lateral ventricle; MD, mediodorsal thalamic nucleus; ML, medial mammillary nucleus,lateral part; MM, medial mammillary nucleus, medial part; mml, medial medullary lamina; mtt, mamillothalamic tract; opt, optic tract; PF, parafascicular thalamic nucleus; PP, posterior parietal cortex; Pu, putamen; Rt, reticular thalamic nucleus; S, subiculum; STG, superior temporal gyrus; STh, subthalamic nucleus; STS, superior temporal sulcus; SuM, supramammillary nucleus; TE, temporal cortex; VL, ventral lateral thalamic nucleus; VPL, ventral posterolateral thalamic nucleus; ZI, zona incerta.






5, 20, 21, 22, 23, 42, 50, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A2, secondary auditory region; ADH, dorsal hypothalamic area; BST, bed nucleus of the stria terminalis; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; CL, central lateral thalamic nucleus; CM, central medial thalamic nucleus; cp, cerebral peduncle,basal part; DG, dentate gyrus; Er, entorhinal cortex; f, fornix; GP, globus pallidus; Hb, habenular nucleus; ic, internal capsule; ITG, inferotemporal gyrus; LD, lateral dorsal thalamic nucleus; If, lateral fissure; LH, lateral hypothalamic area, LV, lateral ventricle; MD, mediodorsal thalamic nucleus; ML, medial mammillary nucleus,lateral part; MM, medial mammillary nucleus,medial part; opt, optic tract; PP, posterior parietal cortex; Pu, putamen; PV, paraventricular thalamic nucleus; Rt, reticular thalamic nucleus; S, subiculum; SMH, supramamillary hypothalamic nucleus; SNc, substantia nigra, compact; SNr, substantia nigra, reticular part; st, stria terminalis; STG, superior temporal gyrus; STh, subthalamic nucleus; STS, superior temporal sulcus; TE, temporal cortex; VL, ventral lateral thalamic nucleus; VPL, ventral posterolateral thalamic nucleus; VPM, ventral posteromedial thalamic nucleus; ZI, zona incerta.






5, 20, 21, 22, 23, 42, 50, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A2, secondary auditory region; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; CL, central lateral thalamic nucleus; CM, central medial thalamic nucleus; cp, cerebral peduncle, basal part: DG, dentate gyrus; Er, entorhinal cortex; f, fornix; Hb, habenular nucleus; ic, internal capsule; ITG, inferotemporal gyrus; LD, lateral dorsal thalamic nucleus; If, lateral fissure; LG, lateral geniculate nucleus; LH, lateral hypothalamic area; LV, lateral ventricle; MD, mediodorsal thalamic nucleus; ML, medial mammillary nucleus, nedial part; opt, optic tract; PF, parafascicular thalamic nucleus; PH, posterior hypothalamic area; PP, posterior parietal cortex; Pu, putamen; PV, paraventricular thalamic nucleus; RO, radiatio optica; Rt, reticular thalamic nucleus; S, subiculum; SN, substantia nigra; STG, superior temporal gyrus; STh, subthalamic nucleus; STS, superior temporal sulcus; TE, temporal cortex; VL, ventral lateral thalamic nucleus; VPL, ventral posterolateral thalamic nucleus; VPM, ventral posteromedial thalamic nucleus; VTA, ventral tegmental area; ZI, zona incerta.







7, 20, 21, 22, 23, 42, 50, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A2, secondary auditory region; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; CL, central lateral thalamic nucleus; CM, central medial thalamic nucleus; cp, cerebral peduncle, basal part; DG, dentate gyrus; Er, entorhinal cortex; EW, Edinger-Westphal nucleus; f, fornix; H, campus Foreli; Hb, habenular nucleus; ic, internal capsule; IP, interpeduncular nucleus; ITG, inferotemporal gyrus; If, lateral fissure; LG, lateral geniculate nucleus; LV, lateral ventricle; MD, mediodorsal thalamic nucleus; PF, parafascicular thalamic nucleus; PP, posterior parietal cortex; Pu, putamen; PV, paraventricular thalamic nucleus; RS, retrosplenial cortex; Rt, reticular thalamic nucleus; S, subiculum; SNc, substantia nigra, compact; SNr, substantia nigra, reticular part; STG, superior temporal gyrus; STS, superior temporal sulcus; TE, temporal cortex; VL, ventral lateral thalamic nucleus; VLG, ventral lateral geniculate nucleus; VPL, ventral posterolateral thalamic nucleus; VPM, ventral posteromedial thalamic nucleus; VTA, ventral tegmental area; ZI, zona incerta.






7, 20, 21, 23, 42, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; A2, secondary auditory region; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; CL, central lateral thalamic nucleus; Cl, claustrum; CM, central medial thalamic nucleus; cp, cerebral peduncle, basal part; DG, dentate gyrus; Er, entorhinal cortex; EW, Edinger-Westphal nucleus; f, fornix; H, campus Foreli; ic, internal capsule; IP, interpeduncular nucleus; ITG, inferotemporal gyrus; LG, lateral geniculate nucleus; LHb, lateral habenular nucleus; LP, lateral posterior nucleus; MD, mediodorsal thalamic nucleus; MHb, medial habenular nucleus; PF, parafascicular thalamic nucleus; PP, posterior parietal cortex; Pu, putamen; PV, paraventricular thalamic nucleus; RO, radiatio optica; RS, retrosplenial cortex; Rt, reticular thalamic nucleus; S, subiculum; SNc, substantia nigra, compact; SNr, substantia nigra, reticular part; STS, superior temporal sulcus; VPL, ventral posterolateral thalamic nucleus; VPM, ventral posteromedial thalamic nucleus; VTA, ventral tegmental area.





7, 20, 21,23, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; AP, anterior pulvinar; Aq, aqueduct; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; CM, central medial thalamic nucleus; cp, cerebral peduncle,basal part; DG, dentate gyrus; eml, external medullary lamina; Er, entorhinal cortex; EW, Edinger-Westphal nucleus; f, fornix; FRT, formatio reticularis tegmenti; GC, substantia grisea centralis; IP, interpeduncular nucleus; ITG, inferotemporal gyrus; LG, lateral geniculate nucleus; LHb, lateral habenular nucleus; LP, lateral posterior nucleus; LV, lateral ventricle; MD, mediodorsal thalamic nucleus; MHb, medial habenular nucleus; opt, optic tract; PF, parafascicular thalamic nucleus; PP, posterior parietal cortex; RS, retrosplenial cortex; Rt, reticular thalamic nucleus; S, subiculum; SNc, substantia nigra, compact; SNr, substantia nigra, reticular part; STS, superior temporal sulcus; VPL, ventral posterolateral thalamic nucleus; VPM, ventral posteromedial thalamic nucleus; VTA, ventral tegmental area.






7, 20, 21, 23, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; CM, central medial thalamic nucleus; cp, cerebral peduncle,basal part; DG, dentate gyrus; eml, external medullary lamina; Er, entorhinal cortex; EW, Edinger-Westphal nucleus; f, fornix; fr, fasciculus retroflexus; FRT, formatio reticularis tegmenti; GC, substantia grisea centralis; IP, interpeduncular nucleus; ITG, inferotemporal gyrus; LG, lateral geniculate nucleus; LHb, lateral habenular nucleus; LP, lateral posterior nucleus; LV, lateral ventricle; MD, mediodorsal thalamic nucleus; MHb, medial habenular nucleus; opt, optic tract; PF, parafascicular thalamic nucleus; PP, posterior parietal cortex; Pp, peripeduncular nucleus; RS, retrosplenial cortex; Rt, reticular thalamic nucleus; S, subiculum; SNc, substantia nigra, compact; SNr, substantia nigra, reticular part; STS, superior temporal sulcus; VPL, ventral posterolateral thalamic nucleus; VTA, ventral tegmental area.


cNissl-40







7, 20, 21, 23, Brodmann's parcellation of the cortex; 3V, 3rd ventricle; Apul, anterior pulvinar; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; CM, central medial thalamic nucleus; cp, cerebral peduncle,basal part; DG, dentate gyrus; Er, entorhinal cortex; EW, Edinger-Westphal nucleus; f, fornix; fr, fasciculus retroflexus; H, campus Foreli; ic, internal capsule; IP, interpeduncular nucleus; IPul, inferior pulvinar; ITG, inferotemporal gyrus; LD, lateral dorsal thalamic nucleus; LG, lateral geniculate nucleus; LHb, lateral habenular nucleus; LPul, lateral pulvinar; LV, lateral ventricle; MD, mediodorsal thalamic nucleus; MG, medial geniculate nucleus; MHb, medial habenular nucleus; MPul, medial pulvinar; PF, parafascicular thalamic nucleus; PP, posterior parietal cortex; Pu, putamen; PV, paraventricular thalamic nucleus; RN, red nucleus; RO, radiatio optica; RS, retrosplenial cortex; Rt, reticular thalamic nucleus; S, subiculum; SNc, substantia nigra, compact; SNr, substantia nigra, reticular part; st, stria terminalis; STS, superior temporal sulcus; VTA, ventral tegmental area.







7, 20, 21, 23, Brodmann's parcellation of the cortex; 3N, oculomotor nucleus; 3V, 3rd ventricle; Apul, anterior pulvinar; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; Cg, cingulate cortex; CM, central medial thalamic nucleus; cp, cerebral peduncle, basal part; DG, dentate gyrus; Er, entorhinal cortex;f, fornix; H, campus Foreli; IP, interpeduncular nucleus; IPul, inferior pulvinar; ITG, inferotemporal gyrus; LG, lateral geniculate nucleus; LHb, lateral habenular nucleus; LPul, lateral pulvinar; LV, lateral ventricle; MD, medial dorsal thalamic nucleus; MG, medial geniculate nucleus; MHb, medial habenular nucleus; MPul, medial pulvinar; MT, Middle temporal visual area; MTc, Middle temporal crescent visual area; PF, parafascicular thalamic nucleus; PP, posterior parietal cortex; RN, red nucleus; RO, radiatio optica; RS, retrosplenial cortex; Rt, reticular thalamic nucleus; S, subiculum; SNc, substantia nigra, compact; SNr, substantia nigra, reticular part; st, stria terminalis; STS, superior temporal sulcus; VTA, ventral tegmental area.






7, 20, 21, 23, Brodmann's parcellation of the cortex; 3N, oculomotor nucleus; 3V, 3rd ventricle; Aq, aqueduct; bsc, brachium of the superior colliculus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; cp, cerebral peduncle, basal part; DG, dentate gyrus; DpMe, deep mesence-phalic nucleus; Er, entorhinal cortex; f, fornix; FRTM, formatio reticularis tegmenti mesencephali; IP, interpeduncular nucleus; IPul, inferior pulvinar; ITG, inferotemporal gyrus; Ifp, longitudinal fasciculus of the pons; LG, lateral geniculate nucleus; LHb, lateral habenular nucleus; LPul, lateral pulvinar; LV, lateral ventricle; mcp, middle cerebellar peduncle; MG, medial geniculate nucleus; MHb, medial habenular nucleus; MPul, medial pulvinar; MT, middle temporal visual area; MTc, Middle temporal crescent visual area; PAG, periaqueductal gray; Pn, pontine nuclei; PP, posterior parietal cortex; RA, radiatio acustica; RN, red nucleus; RS, retrosplenial cortex; Rt, reticular thalamic nucleus; S, subiculum; SC, superior colliculus; SNc, substantia nigra, compact; SNr, substantia nigra, reticular part; STS, superior temporal sulcus; VTA, ventral tegmental area; xscp, decussation of the superior cerebellar peduncle.





7, 20, 21, 23, Brodmann's parcellation of the cortex; 3ND, oculomotor nucleus, dorsal part; 3NV, oculomotor nucleus, ventral part; 3V, 3rd ventricle; Aq, aqueduct; bsc, brachium of the superior colliculus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; cp, cerebral peduncle, basal part; DG, dentate gyrus; DpMe, deep mesencephalic nucleus; eml, external medullary lamina; Er, entorhinal cortex; f, fornix; IP, interpeduncular nucleus; IPul, inferior pulvinar; ITG, inferotemporal gyrus; Ifp, longitudinal fasciculus of the pons; LHb, lateral habenular nucleus; LPul, lateral pulvinar; LV, lateral ventricle; mcp, middle cerebellar peduncle; MG, medial geniculate nucleus; MHb, medial habenular nucleus; ml, medial lemniscus; MPul, medial pulvinar; MT, middle temporal visual area; MTc, middle temporal crescent visual area; PAG, periaqueductal gray; Pn, pontine nuclei; PP, posterior parietal cortex; RA, radiatio acustica; RN, red nucleus; RS, retrosplenial cortex; Rt, reticular thalamic nucleus; S, subiculum; SC, superior colliculus; SNc, substantia nigra, compact; SNr, substantia nigra, reticular part; STS, superior temporal sulcus; xscp, decussation of the superior cerebellar peduncle.






7, 20, 21, 23, Brodmann's parcellation of the cortex; 3ND, oculomotor nucleus, dorsal part; 3NV, oculomotor nucleus, ventral part; Aq, aqueduct; bsc, brachium of the superior colliculus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; DG, dentate gyrus; Er, entorhinal cortex; f, fornix; FRTM, formatio reticularis tegmenti mesencephali; IP, interpeduncular nucleus; IPul, inferior pulvinar; ITG, inferotemporal gyrus; Ifp, longitudinal fasciculus of the pons; LPul, lateral pulvinar; LV, lateral ventricle; mcp, middle cerebellar peduncle; MG, medial geniculate nucleus; ml, medial lemniscus; MPul, medial pulvinar; MT, middle temporal visual area; MTc, middle temporal crescent visual area; PAG, periaqueductal gray; pc, posterior commissure; PP, posterior parietal cortex; RA, radiatio acustica; RN, red nucleus; RS, retrosplenial cortex; Rt, reticular thalamic nucleus; RTP, nucleus reticularis tegmenti pontis; S, subiculum; SC, superior colliculus; SN, substantia nigra; STS, superior temporal sulcus; xscp, decussation of the superior cerebellar peduncle.


cNissl-45






7, 20, 21, 23, Brodmann's parcellation of the cortex; 3ND, oculomotor nucleus, dorsal part; 3NV, oculomotor nucleus, ventral part; Aq, aqueduct; bsc, brachium of the superior colliculus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; csc, commissure of the superior colliculus; DG, dentate gyrus; Er, entorhinal cortex; f, fornix; FRTM, formatio reticularis tegmenti mesencephali; IP, interpeduncular nucleus; IPul, inferior pulvinar; ITG, inferotemporal gyrus; Ifp, longitudinal fasciculus of the pons; LPul, lateral pulvinar; LV, lateral ventricle; mcp, middle cerebellar peduncle; MG, medial geniculate nucleus; ml, medial lemniscus; MPul, medial pulvinar; MT, middle temporal visual area; MTc, middle temporal crescent visual area; PAG, periaqueductal gray; Pn, pontine nuclei; PP, posterior parietal cortex; Pu, putamen; RA, radiatio acustica; RN, red nucleus; RO, radiatio optica; RS, retrosplenial cortex; Rt, reticular thalamic nucleus; S, subiculum; SC, superior colliculus; SNc, substantia nigra, compact; SNr, substantia nigra, reticular part; STS, superior temporal sulcus; xscp, decussation of the superior cerebellar peduncle.






7, 20, 21, 23, Brodmann's parcellation of the cortex; 3N, oculomotor nucleus; Aq, aqueduct; bsc, brachium of the superior colliculus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; CLi, caudal linear nucleus of the raphe; DG, dentate gyrus; DpG, deep gray layer; DpMe; deep mesencephalic nucleus; DR, dorsal raphe nucleus; ECIC, external cortex of the inferior colliculus; Er, entorhinal cortex; f, fornix; InG, intermediate gray layer; IP, interpeduncular nucleus; IPul, inferior pulvinar; ITG, inferotemporal gyrus; Ifp, longitudinal fasciculus of the pons; LPul, lateral pulvinar; LV, lateral ventricle; mcp, middle cerebellar peduncle; MG, medial geniculate nucleus; ml, medial lemniscus; mlf, medial longitudinal fasciculus; MnR, median raphe nucleus; MPul, medial pulvinar; MT, middle temporal visual area; MTc, middle temporal crescent visual area; PAG, periaqueductal gray; pc, posterior commissure; Pn, pontine nuclei; PP, posterior parietal cortex; RS, retrosplenial cortex; S, subiculum; SC, superior colliculus; SCG, superficial gray layer of superior colliculus; STS, superior temporal sulcus; xscp, decussation of the superior cerebellar peduncle.






7, 20, 21, 23, Brodmann's parcellation of the cortex; 4N, trochlear nucleus; Aq, aqueduct; bsc, brachium of the superior colliculus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; csc, commissure of the superior colliculus; DG, dentate gyrus; DR, dorsal raphe nucleus; Er, entorhinal cortex; f, fornix; FRP, formatio reticularis pontis; FRTM; formatio reticularis tegmenti mesencephali; ITG, inferotemporal gyrus; Ifp, longitudinal fasciculus of the pons; LPul, lateral pulvinar; LV, lateral ventricle; mcp, middle cerebellar peduncle; ml, medial lemniscus; mlf, medial longitudinal fasciculus; MPul, medial pulvinar; MT, middle temporal visual area; MTc, middle temporal crescent visual area; NCS, central superior tegmental nucleus; PAG, periaqueductal gray; PP, posterior parietal cortex; RS, retrosplenial cortex; RTP, nucleus reticularis tegmenti pontis; S, subiculum; SC, superior colliculus; STS, superior temporal sulcus; VLP, Ventrolateral posterior extrastriate area; xscp, decussation of the superior cerebellar peduncle.






19, 20, 21, 23, Brodmann's parcellation of the cortex; Aq, aqueduct; bic, brachium of the inferior colliculus; bsc, brachium of the superior colliculus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cc, corpus callosum; Cd, caudate nucleus; DA, dorsoanterior extrastriate area; DG, dentate gyrus; DpMe, deep mesencephalic nucleus; DR, dorsal raphe nucleus; Er, entorhinal cortex; ITG, inferotemporal gyrus; Ifp, longitudinal fasciculus of the pons; II, lateral lemniscus; LV, lateral ventricle; mcp, middle cerebellar peduncle; mI, medial lemniscus; mIf, medial longitudinal fasciculus; MT, middle temporal visual area; MTc, middle temporal crescent visual area; PAG, periaqueductal gray; Pn, pontine nuclei; PrS, presubiculum; PuI, pulvinar nuclei; RS, retrosplenial cortex; S, subiculum; SC, superior colliculus; scp, superior cerebellar peduncle(brachium conjunctivum); vhc, ventral hippocampal commissure; VLP, ventrolateral posterior extrastriate area.

cCalb-49






18, 19, 20, 23, Brodmann's parcellation of the cortex; 4N, trochlear nucleus; Aq, aqueduct; bic, brachium of the inferior colliculus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; DA, dorsoanterior extrastriate area; DG, dentate gyrus; DpMe, deep mesencephalic nucleus; DR, dorsal raphe nucleus; ITG, inferotemporal gyrus; Ifp, longitudinal fasciculus of the pons; LV, lateral ventricle; mcp, middle cerebellar peduncle; ml, medial lemniscus; mlf, medial longitudinal fasciculus; Mo5N, trigeminal motor nucleus; MT, middle temporal visual area; MTc, middle temporal crescent visual area; PAG, periaqueductal gray; Pn, pontine nuclei; RS, retrosplenial cortex; S, subiculum; SC, superior colliculus; V2, secondary visual area; VLA, ventrolateral anterior extrastriate area; VR, ventrolateral posterior extrastriate area; VR, ventral raphe nucleus.


cNissI-50






18, 19, 20, 23, Brodmann's parcellation of the cortex; 4N, trochlear nucleus; Aq, aqueduct; bic, brachium of the inferior colliculus; cic, commissure of the inferior colliculus; DA, dorsoanterior extrastriate area; DR, dorsal raphe nucleus; IC, inferior colliculus; ITG, inferotemporal gyrus; Ifp, longitudinal fasciculus of the pons; II, lateral lemniscus; LV, lateral ventricle; mcp, middle cerebellar peduncle; Me5N, mesencephalic trigeminal nucleus; ml, medial lemniscus; mlf, medial longitudinal fasciculus; Mo5N, trigeminal motor nucleus; MT, middle temporal visual area; MTc, middle temporal crescent visual area; PAG, periaqueductal gray; Pn, pontine nuclei; RS, retrosplenial cortex; S, subiculum; SC, superior colliculus; V2, second visual area; VLA, ventrolateral anterior extrastriate area; VLP, ventrolateral posterior extrastriate area; VR, ventral raphe nucleus.







18, 19, 19m, 20, 23, Brodmann's parcellation of the cortex; 4N, trochlear nucleus; Aq, aqueduct; cic, commissure of the inferior colliculus; DA, dorsoanterior extrastriate area; DR, dorsal raphe nucleus; IC, inferior colliculus; ITG, inferotemporal gyrus; Ifp, longitudinal fasciculus of the pons; II, lateral lemniscus; LV, lateral ventricle; m5/s5, motor root of the trigeminal nerve/sensory root of the trigeminal nerve; mcp, middle cerebellar peduncle; Me5N, mesencephalic trigeminal nucleus; ml, medial lemniscus; mlf, medial longitudinal fasciculus; Mo5N, trigeminal motor nucleus; MTc, middle temporal crescent visual area; PAG, periaqueductal gray; Pn, pontine nuclei; RS, retrosplenial cortex; S, subiculum; SC, superior colliculus; SO, superior olive; V2, second visual area; VLA, ventrolateral anterior extrastriate area; VLP, ventrolateral posterior extrastriate area; VR, ventral raphe nucleus; xscp, decussation of the superior cerebellar peduncle.


cNissI-52






18, 19, 19m, 23, Brodmann's parcellation of the cortex; Aq, aqueduct; cic, commissure of the inferior colliculus; DA, dorsoanterior extrastriate area; DR, dorsal raphe nucleus; IC, inferior colliculus; Ifp, longitudinal fasciculus of the pons; II, lateral lemniscus; LV, lateral ventricle; mcp, middle cerebellar peduncle; ml, medial lemniscus; mlf, medial longitudinal fasciculus; Mo5N, trigeminal motor nucleus; MTc, middle temporal crescent visual area; PAG, periaqueductal gray; Pn, pontine nuclei; S, subiculum; SC, superior colliculus; scp, superior cerebellar peduncle(brachium conjunctivum); SO, superior olive; V2, second visual area; VLA, ventrolateral anterior extrastriate area; VLP, ventrolateral posterior extrastriate area; VR, ventral raphe nucleus.






4V, 4th ventricle; 6N, abducens nucleus; 6n, root of abducens nerve; 7n, facial nerve; Cb1, cerebellar lobule 1; Cb2, cerebellar lobule 2; Cb3, cerebellar lobule 3; Cb4, cerebellar lobule 4; Cb5; cerebellar lobule 5; Cb6, cerebellar lobule 6; g7, genu of the facial nerve; LC, locus coeruleus; mcp, middle cerebellar peduncle; ml, medial lemniscus; mlf, medial longitudinal fasciculus; Mo5N, motor trigeminal nucleus; Pn, pontine nuclei; PPFI, posterior paraflocculus; py, pyramidal tract; RMg, nucleus raphe magnus; scp, superior cerebellar peduncle (brachium conjunctivum); SO, superior olive; Sp5, spinal trigeminal nucleus; SVeN, superior vestibular nucleus.






4V, 4th ventricle; 6N, abducens nucleus; 6n, root of abducens nerve; 7n, facial nerve; 8n, vestibulocochlear nerve; Cb1, cerebellar lobule 1; Cb2, cerebellar lobule 2; Cb3, cerebellar lobule 3; Cb4, cerebellar lobule 4; Cb5; cerebellar lobule 5; Cb6, cerebellar lobule 6; g7, genu of the facial nerve; icp, inferior cerebellar peduncle (restiform body); LVeN, lateral vestibular nucleus; mcp, middle cerebellar peduncle; ml, medial lemniscus; mlf, medial longitudinal fasciculus; MVeN, medial vestibular nucleus; Pn, pontine nuclei; PPFI, posterior paraflocculus; py, pyramidal tract; RMg, nucleus raphe magnus; ROb, raphe obscurus nucleus; scp, superior cerebellar peduncle (brachium conjunctivum); SO, superior olive; Sp5, spinal trigeminal nucleus; sp5, spinal trigeminal tract; SVeN, superior vestibular nucleus; VCA, ventral cochlear nucleus, anterior part; vsc, ventral spinocerebellar tract.






4V, 4th ventricle; 7N, facial nucleus; 8n, vestibulocochlear nerve; Cb3, cerebellar lobule 3; Cb4, cerebellar lobule 4; Cb5; cerebellar lobule 5; Cb6, cerebellar lobule 6; g7, genu of the facial nerve; icp, inferior cerebellar peduncle (restiform body); LPB, lateral parabrachial nucleus; mcp, middle cerebellar peduncle; ml, medial lemniscus; mlf, medial longitudinal fasciculus; MPB, medial parabrachial nucleus; PPFI, posterior paraflocculus; Pr, prepositus nucleus; py, pyramidal tract; RP, nucleus raphe pallidus; RTP, nucleus reticularis tegmenti pontis; scp, superior cerebellar peduncle (brachium conjunctivum); Sp5, spinal trigeminal nucleus; sp5, spinal trigeminal tract; VCA, ventral cochlear nucleus, anterior part; VeN, vestibular nuclei; vsc, ventral spinocerebellar tract.






4V, 4th ventricle; 7N, facial nucleus; 8n, vestibulocochlear nerve; Cb3, cerebellar lobule 3; Cb4, cerebellar lobule 4; Cb5; cerebellar lobule 5; Cb6, cerebellar lobule 6; icp, inferior cerebellar peduncle (restiform body); LPB, lateral parabrachial nucleus; LVeN, lateral vestibular nucleus; mcp, middle cerebellar peduncle; mlf, medial longitudinal fasciculus; MPB, medial parabrachial nucleus; MVeN, medial vestibular nucleus; Pn, pontine nuclei; PPFI, posterior paraflocculus; py, pyramidal tract; RMg, nucleus raphe magnus; ROb, raphe obscurus nucleus; scp, superior cerebellar peduncle (brachium conjunctivum); Sp5, spinal trigeminal nucleus; sp5, spinal trigeminal tract; VCA, ventral cochlear nucleus, anterior part.





4V, 4th ventricle; 7n, facial nerve; 7N, facial nucleus; 8n, vestibulocochlear nerve; Cb4, cerebellar lobule 4; Cb5, cerebellar lobule 5; Cb6, cerebellar lobule 6; icp, inferior cerebellar peduncle (restiform body); Int, interposed cerebellar nucleus; IO, inferior olive; Lat, lateral (dentate) cerebellar nucleus; LVeN, lateral vestibular nucleus; mcp, middle cerebellar peduncle; MDR, medullary reticular nucleus; Med, medial (fastigial) cerebellar nucleus; ml, medial lemniscus; mlf, medial longitudinal fasciculus; MVeN, medial vestibular nucleus; PPFI, posterior paraflocculus; Pr, prepositus nucleus; py, pyramidal tract; RMg, nucleus raphe magnus; Sp5, spinal trigeminal nucleus; sp5, spinal trigeminal tract; SVeN, superior vestibular nucleus; VCA, ventral cochlear nucleus, anterior part.






4V, 4th ventricle; 4VLR, lateral recess of the 4th ventricle; 7N, facial nucleus; Cb5, cerebellar lobule 5; Cb6, cerebellar lobule 6; Gi, gigantocellular reticular nucleus; icp, inferior cerebellar peduncle (restiform body); IntA, interposed cerebellar nucleus, anterior part, IntP, interposed cerebellar nucleus, posterior part, IO, inferior olive; Irt, intermediate reticular nucleus; Lat, lateral (dentate) cerebellar nucleus; LVeN, lateral vestibular nucleus; Med, medial (fastigial) cerebellar nucleus; mlf, medial longitudinal fasciculus; MVeNMC, medial vestibular nucleus, magnocellular; MVeNPC, medial vestibular nucleus, parvicellular; PPFI, posterior paraflocculus; Pr, prepositus nucleus; py, pyramidal tract; Sp5, spinal trigeminal nucleus; sp5, spinal trigeminal tract; VCP, ventral cochlear nucleus, posterior part.


cNissI-59







4V, 4th ventricle; Cb5, cerebellar lobule 5; Cb6, cerebellar lobule 6; Cb10, cerebellar lobule 10; CR, central reticular nucleus; DM10, nucleus dorsalis motorius nervi vagi; icp, inferior cerebellar peduncle (restiform body); IntA, interposed cerebellar nucleus, anterior part, IntP, interposed cerebellar nucleus, posterior part, IO, inferior olive; Lat, lateral (dentate) cerebellar nucleus; LRt, lateral reticular nucleus; MDR, medullary reticular nucleus; Med, medial (fastigial) cerebellar nucleus; ml, medial lemniscus; mlf, medial longitudinal fasciculus; PPFI, posterior paraflocculus; Pr, prepositus nucleus; Py, pyramidal tract; RMg, nucleus raphe magnus; Sp5, spinal trigeminal nucleus; sp5, spinal trigeminal tract; VCP, ventral cochlear nucleus, posterior part; VeN, vestibular nuclei.






10N, dorsal motor nucleus of vagus nerve; 10n, vagus nerve; 12N, hypoglossal nucleus; Amb, ambiguus nucleus; Cb5, cerebellar lobule 5; Cb6, cerebellar lobule 6; Cb10, cerebellar lobule 10; DPFI, dorsal paraflocculus; FI, flocculus; icp, inferior cerebellar peduncle (restiform body); Int, interposed cerebellar nucleus; IO, inferior olive; Lat, lateral (dentate) cerebellar nucleus; LRt, lateral reticular nucleus; Med, medial (fastigial) cerebellar nucleus; mI, medial lemniscus; mIf, medial longitudinal fasciculus; py, pyramidal tract; RMg, nucleus raphe magnus; Sim, simple lobule; Sol, solitary nucleus; Sp5, spinal trigeminal nucleus; VeN, vestibular nuclei.







12n, root of hypoglossal nerve; Amb, ambiguus nucleus; Cb5, cerebellar lobule 5; Cb6, cerebellar lobule 6; Cb9, cerebellar lobule 9; DPFI, dorsal paraflocculus; FI, flocculus; icp, inferior cerebellar peduncle (restiform body); Int, interposed cerebellar nucleus; IO, inferior olive; Lat, lateral (dentate) cerebellar nucleus; LRt, lateral reticular nucleus; MDR, medullary reticular nucleus; Med, medial (fastigial) cerebellar nucleus; mI, medial lemniscus; mIf, medial longitudinal fasciculus; PPFI, posterior paraflocculus; Pr, prepositus nucleus; py, pyramidal tract; RP, nucleus raphe pallidus; Sol; solitary nucleus; Sp5, spinal trigeminal nucleus; sp5, spinal trigeminal tract; VeN, vestibular nuclei.






10N, dorsal motor nucleus of vagus nerve; 10n, vagus nerve; 12N, hypoglossal nucleus; 12n, root of hypoglossal nerve; Amb, ambiguus nucleus; Cb5, cerebellar lobule 5; Cb6, cerebellar lobule 6; Cb8, cerebellar lobule 8; Cb9, cerebellar lobule 9; DPFI, dorsal paraflocculus; FI, flocculus; icp, inferior cerebellar peduncle (restiform body); Int, interposed cerebellar nucleus; IO, inferior olive; Lat, lateral (dentate) cerebellar nucleus; LRt, lateral reticular nucleus; Med, medial (fastigial) cerebellar nucleus; mlf, medial longitudinal fasciculus; PPFI, posterior paraflocculus; py, pyramidal tract; RP, nucleus raphe pallidus; Sim, simple lobule; Sol, solitary nucleus; Sp5, spinal trigeminal nucleus; sp5, spinal trigeminal tract; VeN, vestibular nuclei.






10N, dorsal motor nucleus of vagus nerve; ans, ansiform lobule of the cerebellum; Cb5, cerebellar lobule 5; Cb6, cerebellar lobule 6; Cb7, cerebellar lobule 7; Cb8, cerebellar lobule 8; Cb9, cerebellar lobule 9; Cop, copula of the pyramis; CR, central reticular nucleus; Crus1, crus 1 of the ansiform lobule; Crus2, crus 2 of the ansiform lobule; Cu, cuneate nucleus; ECu, external cuneate nucleus; Gr, gracile nucleus; LRt, lateral reticular nucleus; py, pyramidal tract; pyx, pyramidal decussation; Sp5, spinal trigeminal nucleus; sp5, spinal trigeminal tract.

cNissl-64



cTr-65




17, 18, 19, Brodmann's parcellation of the cortex; cal, calcarine sulcus; DA, dorsoanterior extrastriate area; DI, dorsointermediate visual area; DM, dorsomedial visual area; POm, parietooccipital medial area; V1, primary visual cortex; V2, second visual area; VLA, ventrolateral anterior extrastriate area; VLP, ventrolateral posterior extrastriate area.

cNissl-65





cTr-66



17, 18, 19, Brodmann's parcellation of the cortex; cal, calcarine sulcus; DA, dorsoanterior extrastriate area; DI, dorsointermediate visual area; DM, dorsomedial visual area; POm, parietooccipital medial area; V1, primary visual cortex; V2, second visual area; VLA, ventrolateral anterior extrastriate area; VLP, ventrolateral posterior extrastriate area.


cNissl-66






optic tract; PH, posterior hypothalamic area; Pn, pontine nuclei; POA, preoptic area; POm, parietooccipital medial area; Pul, pulvinar nuclei; py, pyramidal tract; pyrs, ventromedial hypothalamic nucleus; VTA, ventral tegmental area; xml, decussation of medial lemniscus; xscp, decussation of the superior cerebellar peduncle; ZI, zona 3a, 3b, 4, 6dr, 6m, 7, 8B, 9, 10, 14, 17, 18, 19, 25, Brodmann's parcellation of the cortex; 3N, oculomotor nucleus; 3V, 3rd ventricle; 4N, trochlear nucleus; 12N, hypoglosmediodorsal thalamic nucleus; ml, medial lemniscus; mlf, medial longitudinal fasciculus; MM, medial mammillary nucleus, medial part; mtt, mamillothalamic tract; opt, sal nucleus; ac, anterior commissure; AM, anteromedial thalamic nucleus; BST, bed nucleus of the stria terminalis; Cb1-Cb10, cerebellar lobule 1-10; cc, corpus callosum; CG, central gray; DA, dorsoanterior extrastriate area; dlf, dorsal longitudinal fasciculus; DM, dorsomedial hypothalamic nucleus; DM (visual area), dorsomedial area; f fornix; Gr. gracile nucleus; Hb, habenular nucleus; IC, inferior colliculus; InfS, infundibular stem; LSD, lateral septal nucleus, dorsal part; M1, primary motor area; MD, pyramis; RN, red nucleus; S1, primary somatosensory area; SC, superior colliculus; SON, supraoptic nucleus; V1, primary visual cortex; V2, second visual area; VMH,


psNissI-01




medial thalamic nucleus; cal, calcarine sulcus; Cb1-Cb10, cerebellar lobule 1-10; cc, corpus callosum; Cd, caudate nucleus; CG, central gray; CM, central medial thaopt, optic tract; Pav, paraventricular hypothalamic nucleus; PH, posterior hypothalamic area; Pn, pontine nuclei; POA, preoptic area; POm, parietooccipital medial area; PON, preoptic nucleus; Pul, pulvinar nuclei; pyrs, pyramis; pyx, pyramidal decussation; RN, red nucleus; S1, primary somatosensory area; SC, superior colliculus; Sep, Septum; SON, supraoptic nucleus; V1, primary visual cortex; V2, second visual area; VA, ventral anterior thalamic nucleus; VMH, ventromedial hypothalamic nucleus; 3a, 3b, 4, 6dr, 6m, 7, 8B, 9, 10, 14, 17, 18, 19, Brodmann's parcellation of the cortex; 4V, 4th ventricle; 12N, hypoglossal nucleus; ac, anterior commissure; AM, anterolamic nucleus; DA, dorsoanterior extrastriate area; DM, dorsomedial hypothalamic nucleus; DM (visual area), dorsomedial area; f, fornix, fr, fasciculus retroflexus; Gr, gracile nucleus; gr, gracile fasciculus; H, campus Foreli; Hb, habenular nucleus; IC, inferior colliculus; InfS, infundibular stem; IsC, Island of Calleja; LC, locus coeruleus; medial lemniscus; mlf, medial longitudinal fasciculus; MM, medial mammillary nucleus, medial part; Mo5N, motor nucleus of trigeminal nerve; mtt, mamillothalamic tract; LV, lateral ventricle; M1, primary motor area; MD, mediodorsal thalamic nucleus; Me5N, mesencephalic trigeminal nucleus; Med, medial (fastigial) cerebellar nucleus; ml, VTA, ventral tegmental area; xscp, decussation of the superior cerebellar peduncle.




aqueductal gray; Pav, paraventricular hypothalamic nucleus; PH, posterior hypothalamic area; Pn, pontine nuclei; POm, parietooccipital medial area; PON, preoptic anterodorsal nucleus; Arc, arcuate hypothalamic nucleus; cal, calcarine sulcus; Cb1-Cb10, cerebellar lobule 1-10; cc, corpus callosum; Cd, caudate nucleus; CM, central medial thalamic nucleus; cp, cerebral peduncle, basal part; Cu, cuneate nucleus; DA, dorsoanterior extrastriate area; DM, dorsomedial hypothalamic nucleus; DM (visual area), dorsomedial area; f, fornix; flm, fasciculus longitudinalis medialis; fr, fasciculus retroflexus; Gi, gigantocellular reticular nucleus; GP, globus pallidus; Gr, gracile nucleus; Hb, habenular nucleus; HDB, nucleus of the horizontal limb of the diagonal band; IC, inferior colliculus; ic, internal capsule; IO, inferior olive; ISC, Island of Calleja; lo, lateral olfactory tract; M1, primary motor area; MD, mediodorsal thalamic nucleus; Me5N, mesencephalic trigeminal nucleus; Med, medial (fastigial) cerebellar nucleus; ml, medial lemniscus; MM, medial mammillary nucleus, medial part; mtt, mamillothalamic tract; MVeN, medial vestibular nucleus; ox, optic chiasm; PAG, perinucleus; Pu, putamen; Pul, pulvinar nuclei; py, pyramidal tract; R, raphe nucleus; RN, red nucleus; S1, primary somatosensory area; SC, superior colliculus; Sep, Sep-3a, 3b, 4, 6d, 6dr, 7, 8Ad, 9, 10, 11, 13, 17, 18, 19, Brodmann's parcellation of the cortex; 6N, abducens nucleus; 11N, accessory nucleus; ac, anterior commissure; AD, tum; SN, substantia nigra; Sol, solitary nucleus; SON, supraoptic nucleus; V1, primary visual cortex; V2, Second visual area; VA, ventral anterior thalamic nucleus; VH. ventral horn; VMH, ventromedial hypothalamic nucleus; VTA, ventral tegmental area.






CM, central medial thalamic nucleus; DA, dorsoanterior extrastriate area; DM (visual area), dorsomedial area; EGP, external globus pallidus; f, fornix; Gr, gracile nucleus; tibular nucleus; ox, optic chiasm; PAG, periaqueductal gray; Pav, paraventricular hypothalamic nucleus; Pn, pontine nuclei; POA, preoptic area; POm, parietooccipital medial area; Pp, peripeduncular nucleus; Pu, putamen; Pul, pulvinar nuclei; py, pyramidal tract; RN, red nucleus; S1, primary somatosensory area; SC, superior colliculus; scp, superior cerebellar peduncle (brachium conjunctivum); SN, substantia nigra; Sol, solitary nucleus; SON, supraoptic nucleus; st, stria terminalis; V1, primary vi-3a, 3b, 4, 6, 7, 8Ad, 9, 10, 11, 13, 17, 18, 19, 46, Brodmann's parcellation of the cortex; 3n, oculomotor nerve or its root; 6N, abducens nucleus; ac, anterior commissure; AD, anterodorsal nucleus; BST, bed nucleus of the stria terminalis; cal, calcarine sulcus; Cb1-Cb10, cerebellar lobule 1-10; cc, corpus callosum; Cd, caudate nucleus; gr, gracile fasciculus; H, campus Foreli; IC, inferior colliculus; IGP, internal globus pallidus; IO, inferior olive; LC, locus coeruleus; LD, lateral dorsal thalamic nucleus; M1, primary motor area; MD, mediodorsal thalamic nucleus; Me5N, mesencephalic trigeminal nucleus; Med, medial (fastigial) cerebellar nucleus; ML, medial mammillary nucleus, lateral part; ml. medial lemniscus; mlf, medial longitudinal fasciculus; Mo5N, motor nucleus of trigeminal nerve; mtt, mamillothalamic tract; MVeN, medial vessual cortex; V2, second visual area; VA, ventral anterior thalamic nucleus; VLM, ventral lateral thalamic nucleus; medial part; VMH, ventromedial hypothalamic nucleus; VPM, ventral posteromedial thalamic nucleus; VTA, ventral tegmental area.




pontine nuclei; PP, posterior parietal cortex; Pu, putamen; Pul, pulvinar nuclei; py, pyramidal tract; RN, red nucleus; Rt, reticular thalamic nucleus; S1, primary somatosensory area; SC, superior colliculus; scp, superior cerebellar peduncle (brachium conjunctivum); SI, substantia innominata; SNc, substantia nigra, compact; SNr, subdorsoanterior extrastriate area; DM (visual area), dorsomedial area; f, fornix; GP, globus pallidus; Gr, gracile nucleus; IC, inferior colliculus; ic, internal capsule; IO, inferior olive; LD, lateral dorsal thalamic nucleus; M1, primary motor area; MD, mediodorsal thalamic nucleus; Med, medial (fastigial) cerebellar nucleus; ml, medial lemnisstantia nigra, reticular part; SON, supraoptic nucleus; V1, primary visual cortex; V2, second visual area; VA, ventral anterior thalamic nucleus; VH, ventral horn; VL, cus; mlf, medial longitudinal fasciculus; MVeN, medial vestibular nucleus; ox, optic chiasm; PAG, periaqueductal gray; Pav, paraventricular hypothalamic nucleus; Pn, 3a; 3b; 4, 6, 7; 8Ad; 9; 10; 17; 18; 19, 46; Brodmann's parcellation of the cortex; ac, anterior commissure; AD, anterodorsal nucleus; cal, calcarine sulcus; Cb1-Cb10 cerebellar lobule 1-10; cc, corpus callosum; Cd, caudate nucleus; CM, central medial thalamic nucleus; cp, cerebral peduncle, basal part; Cu, cuneate nucleus; DA, ventral lateral thalamic nucleus; VLM, ventral lateral thalamic nucleus, medial part; VPM, ventral posteromedial thalamic nucleus; ZI, zona incerta






3a, 3b, 4, 6dc, 6dr, 7, 8Ad, 9, 10, 17, 18, 19, 46, Brodmann's parcellation of the cortex; 5N, trigeminal nucleus; 7N, facial nucleus; ac, anterior commissure; CA1, field us; cp, cerebral peduncle, basal part; DA, dorsoanterior extrastriate area; DG, dentate gyrus; DM (visual area), dorsomedial area; EGP, external globus pallidus; f, fornix; duncle (brachium conjunctivum); SI, substantia innominata; SNc, substantia nigra, compact; SNr, substantia nigra, reticular part; STh, subthalamic nucleus; V1, primary CA1 of hippocampus; CA3, field CA3 of hippocampus; Cb1-Cb10, cerebellar lobule 1-10; cc, corpus callosum; Cd, caudate nucleus; CM, central medial thalamic nuclereticular thalamic nucleus; RTP, nucleus reticularis tegmenti pontis; S, subiculum; S1, primary somatosensory area; SC, superior colliculus; scp, superior cerebellar peortex; V2, Second visual area; VA, ventral anterior thalamic nucleus; VACo, ventral anterior cortical nucleus of the amygdale; VL, ventral lateral thalamic nucleus; IC, inferior colliculus; ic, internal capsule; icp, inferior cerebellar peduncle (restiform body); IGP, internal globus pallidus; Lat, lateral (dentate) cerebellar nucleus; LD, lateral dorsal thalamic nucleus; M1, primary motor area; MD, mediodorsal thalamic nucleus; ml, medial lemniscus; opt, optic tract; OF, orbitofrontal cortex; PAG, periaqueductal gray; Pn, pontine nuclei; PP, posterior parietal cortex; Pp, peripeduncular nucleus; Pu, putamen; Pul, pulvinar nuclei; py, pyramidal tract; RN, red nucleus; Rt, VPM, ventral posteromedial thalamic nucleus; ZI, zona incerta.

psNissI-06





3a, 3b, 4, 6dc, 6dr, 7, 8Av, 9, 10, 17, 18, 19, Brodmann's parcellation of the cortex; 5N, trigeminal nucleus; 7N, facial nucleus; AA, anterior amygdaloid area; ac, anterior commissure; ans, ansiform lobule of the cerebellum; BM, basomedial amygdaloid nucleus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cal, calcarine mary somatosensory area; SI, substantia innominata; Sim, simple lobule; SN, substantia nigra; SON, supraoptic nucleus; STh, subthalamic nucleus; V1, primary visual cp, cerebral peduncle, basal part; Crus 1, crus 1 of the ansiform lobule; Crus 2 of the ansiform lobule; DA, dorsoanterior extrastriate area; DG, dentate gyrus; DM visual area), dorsomedial area; EGP, external globus pallidus; f, fornix; IC, inferior colliculus; ic, internal capsule; icp, inferior cerebellar peduncle (restiform body); IGP, nternal globus pallidus; Lat, lateral (dentate) cerebellar nucleus; LD, lateral dorsal thalamic nucleus; LVeN, lateral vestibular nucleus; mcp, middle cerebellar peduncle; paramedian lobule; PP, posterior parietal cortex; PPFI, posterior paraflocculus; Pu, putamen; Pul, pulvinar nuclei; Rt, reticular thalamic nucleus; S, subiculum; S1, prisulcus; Cb5, cerebellar lobule 5; Cb6, cerebellar lobule 6; cc, corpus callosum; Cd, caudate nucleus; CM, central medial thalamic nucleus; Cop, copula of the pyramis; M1, primary motor area; MD, mediodorsal thalamic nucleus; Me, medial amygdaloid nucleus; MG, medial geniculate nucleus; OF, orbitofrontal cortex; opt, optic tract; PM, cortex; V2, second visual area; VA, ventral anterior thalamic nucleus; VL, ventral lateral thalamic nucleus; VPFI, ventral paraflocculus; ZI, zona incerta.






3a, 3b, 4, 6dc, 6dr, 7, 8Ad, 10, 17, 18, 19, 46. Brodmann's parcellation of the cortex; AA, anterior amygdaloid area; ac, anterior commissure; ans, ansiform lobule of the cerebellum; BL, basolateral amygdaloid nucleus; BM, basomedial amygdaloid nucleus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; Cd, caudate nucleus; Ce, central amygdaloid nucleus; Cop, copula of the pyramis; cp, cerebral peduncle, basal part; Crus1, crus 1 of the ansiform lobule; Crus2, crus 2 of the ansiform subiculum; S1, primary somatosensory area; Sim, simple lobule; SON, supraoptic nucleus; V1, primary visual cortex; V2, second visual area; VCA, ventral cochlear orbitofrontal cortex; opt, optic tract; Per, periamygdaloid nucleus; PM, paramedian lobule; PPFI, posterior paraflocculus; Pu, putamen; Rt, reticular thalamic nucleus; S, lobule; DA, dorsoanterior extrastriate area; DG, dentate gyrus; DM (visual area), dorsomedial area; EGP, external globus pallidus; Er, entorhinal cortex; ic, internal capsule; IGP, internal globus pallidus; IPul, inferior pulvinar; LP, lateral posterior nucleus; LPul, lateral pulvinar; M1, primary motor area; MG, medial geniculate nucleus; OF, nucleus, anterior part; VLP, Ventrolateral posterior extrastriate area; VPFI, ventral paraflocculus; VPL, ventral posterolateral thalamic nucleus; ZI, zona incerta.






3a, 3b, 4, 6dc, 7, 8Av, 12/45, 17, 18, 46, Brodmann's parcellation of the cortex, AA, anterior amygdaloid area; ac, anterior commissure; BL, basolateral amygdaloid nucleus; BM, basomedial amygdaloid nucleus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cal, calcarine sulcus; cc, corpus callosum; Cd, caudate capsule; IGP, internal globus pallidus; LG, lateral geniculate nucleus; LV, lateral ventricle; M1, primary motor area; Per, periamygdaloid nucleus; Pu, putamen; Pul, pulvinar nuclei; S, subiculum; S1, primary somatosensory area; S1, substantia innominata; V1, primary visual cortex; V2, second visual area; VLP, ventrolateral posterior exnucleus; Ce, central amygdaloid nucleus; DG, dentate gyrus; DM (visual area), dorsomedial area; EGP, external globus pallidus; Er, entorhinal cortex; f, fornix; ic, internal trastriate area.




3a, 3b, 4/6, 7, 8Av, 12/45, 17, 18, 46, Brodmann's parcellation of the cortex; AA, anterior amygdaloid area; ac, anterior commissure; BL, basolateral amygdaloid nucleus; BM, basomedial amygdaloid nucleus; CA1, field CA1 of hippocampus; CA3, field CA3 of hippocampus; cal, calcarine sulcus; cc, corpus callosum; Cd, caudate nucleus; Ce, central amygdaloid nucleus; DG, dentate gyrus; EGP, external globus pallidus; Er, entorhinal cortex; f, fornix; ic, internal capsule; La, lateral amygdaloid nucleus; LG, lateral geniculate nucleus; LV, lateral ventricle; Per, periamygdaloid nucleus; Pu, putamen; Pul, pulvinar nuclei; S1, primary somatosensory area; SI, substantia innominata; st, stria terminalis; V1, primary visual cortex; V2, Second visual area.






CA3, field CA3 of hippocampus; cal, calcarine sulcus; Cd, caudate nucleus; CeL, central amygdaloid nucleus, lateral division; DG, dentate gyrus; EGP, external globus pallidus; Er, entorhinal cortex; ic, internal capsule; La, lateral amygdaloid nucleus; LG, lateral geniculate nucleus; Per, periamygdaloid nucleus; Pu, putamen; Pul, pulvinar nuclei; Rt, reticular thalamic nucleus; st, stria terminalis; S1, primary somatosensory area; V1, primary visual cortex; V2, second visual area; Z1, zona incerta. 3a, 3b, 4/6, 5, 7, 8Av, 12/45, 17, 18, 19, Brodmann's parcellation of the cortex; AA, anterior amygdaloid area; ac, anterior commissure; CA1, field CA1 of hippocampus;






3a, 3b, 4/6, 5, 7, 8Av, 12/45, 17, 18, 19, 41, Brodmann's parcellation of the cortex; A1, primary auditory area; AA, anterior amygdaloid area; ac, anterior commissure; CA1, field CA1 of hippocampus; cal, calcarine sulcus; cc, corpus callosum; Cd, caudate nucleus; CeL, central amygdaloid nucleus, lateral division; DG, dentate gyrus; EGP, external globus pallidus; Er, entorhinal cortex; f, fornix; ic, internal capsule; La, lateral amygdaloid nucleus; If, lateral fissure; LG, lateral geniculate nucleus; LV, lateral ventricle; Pu, putamen; Pul, pulvinar nuclei; S1, primary somatosensory area; st, stria terminalis; V1, primary visual cortex; V2, second visual area.



3a, 3b, 17, 18, 41, Brodmann's parcellation of the cortex; A1, primary auditory area; CA1, field CA1 of hippocampus; Cln, insularis cortex; Cl, claustrum; ec, external capsule; Er, entorhinal cortex; Fr, frontal cortex; ic, internal capsule; If, lateral fissure; LV, lateral ventricle; MT, middle temporal area; MTc, middle temporal cortex; PC, second visual area.



17, 18, 41, Brodmann's parcellation of the cortex; A1, primary auditory area; CA1, field CA1 of hippocampus; Cd, caudate nucleus; Cln, insularis cortex; Cl, claustrum; ec, external capsule; Er, entorhinal cortex; Fr, frontal cortex; ic, internal capsule; If, lateral fissure; LV, lateral ventricle; MT, middle temporal visual area; MTc, middle temporal sulcus; TE, temporal cortex; V1, primary visual cortex; V2, second visual area.



6. Literatures on the Anatomy of Marmoset Brain

Neocortex (general)

- Brysch I, Brysch W, Creutzfeldt O, Hayes NL, Schlingensiepen KH (1984) The second, intralaminar thalamocortical projection system. Anat Embryol (Berl). 169(2):111-8
- Brysch W, Brysch I, Creutzfeldt OD, Schlingensiepen R, Schlingensiepen KH (1990) The topology of the thalamocortical projections in the marmoset monkey (Callithrix jacchus). Exp Brain Res. 81(1):1-17
- Burish MJ, Stepniewska I, Kaas JH (2008) Microstimulation and architectonics of frontoparietal cortex in common marmosets (Callithrix jacchus). J Comp Neurol. 507(2):1151-68
- Cappe C, Barone P (2005) Heteromodal connections supporting multisensory integration at low levels of cortical processing in the monkey. Eur J Neurosci. 22(11):2886-902
- Collins CE, Lyon DC, Kaas JH (2005) Distribution across cortical areas of neurons projecting to the superior colliculus in new world monkeys. Anat Rec A Discov Mol Cell Evol Biol. 285(1):619-27
- Collins P, Wilkinson LS, Everitt BJ, Robbins TW, Roberts AC (2000) The effect of dopamine depletion from the caudate nucleus of the common marmoset (Callithrix jacchus) on tests of prefrontal cognitive function. Behav Neurosci. 114(1):3-17
- Dias R. Robbins TW, Roberts AC (1996) Primate analogue of the Wisconsin Card Sorting Test: effects of excitotoxic lesions of the prefrontal cortex in the marmoset. Behav Neurosci. 110(5):872-86
- Dick A, Kaske A, Creutzfeldt OD (1991) Topographical and topological organization of the thalamocortical projection to the striate and prestriate cortex in the marmoset (Callithrix jacchus), Exp Brain Res. 84(2):233-53
- Divac I, Marinković S, Mogensen J, Schwerdtfeger W, Regidor J (1987) Vertical ascending connections in the isocortex. Anat Embryol (Berl). 175(4):443-55
- Elston GN (2003) The pyramidal neuron in occipital, temporal and prefrontal cortex of the owl monkey (Aotus trivirgatus): regional specialization in cell structure. Eur J Neurosci. 17(6):1313-8
- 11. Elston GN, Benavides-Piccione R, DeFelipe J (2001) The pyramidal cell in cognition: a comparative study in human and monkey. J Neurosci. 21(17):RC163
- Elston GN, Benavides-Piccione R, Elston A, Zietsch B, Defelipe J, Manger P, Casagrande V, Kaas JH (2006) Specializations of the granular prefrontal cortex of primates: implications for cognitive processing. Anat Rec A Discov Mol Cell Evol Biol. 288(1):26-35
- Elston GN, Elston A, Casagrande V, Kaas JH (2005) Pyramidal neurons of granular prefrontal cortex of the galago: complexity in evolution of the psychic cell in primates. Anat Rec A Discov Mol Cell Evol Biol. 285(1):610-8
- Everitt BJ, Sirkiä TE, Roberts AC, Jones GH, Robbins TW (1988) Distribution and some projections of cholinergic neurons in the brain of the common marmoset, Callithrix jacchus. J Comp Neurol. 271(4):533-58
- Ferris CF, Snowdon CT, King JA, Sullivan JM Jr, Ziegler TE, Olson DP, Schultz-Darken NJ, Tannenbaum PL, Ludwig R, Wu Z, Einspanier A, Vaughan JT, Duong TQ (2004) Activation of neural pathways associated with sexual arousal in non-human primates. J Magn Reson Imaging. 19(2):168-75
- Fine A, Hoyle C, Maclean CJ, Levatte TL, Baker HF, Ridley RM (1997) Learning impairments following injection of a selective cholinergic immunotoxin, ME20.4 IgG-saporin, into the basal nucleus of Meynert in monkeys. Neuroscience. 81(2):331-43
- Fonta C, Négyessy L, Renaud L, Barone P (2004) Areal and subcellular localization of the ubiquitous alkaline phosphatase in the primate cerebral cortex: evidence for a role in neurotransmission. Cereb Cortex, 14(6):595-609
- Fonta C, Negyessy L, Renaud L, Barone P (2005) Postnatal development of alkaline phosphatase activity correlates with the maturation of neurotransmission in the cerebral cortex. J Comp Neurol. 486(2):179-96
- Gebhard R, Zilles K, Schleicher A, Everitt BJ, Robbins TW, Divac I (1995) Parcellation of the frontal cortex of the New World monkey Callithrix jacchus by eight neurotransmitter-binding sites. Anat Embryol (Berl). 191(6):509-17
- Geula C, Nagykery N, Wu CK (2002) Amyloid-beta deposits in the cerebral cortex of the aged common marmoset (Callithrix jacchus): incidence and chemical composition. Acta Neuropathol. 103(1):48-58
- 21. Guldin WO, Mirring S, Grüsser OJ (1993) Connections from the neocortex to the vestibular brain stem nuclei in the common marmoset. Neuroreport. 5(2):113-6
- Heilbroner PL, Holloway RL (1988) Anatomical brain asymmetries in New World and Old World monkeys: stages
 of temporal lobe development in primate evolution. Am J Phys Anthropol. 76(1):39-48
- Hornung JP, Celio MR (1992) The selective innervation by serotoninergic axons of calbindin-containing interneurons in the neocortex and hippocampus of the marmoset. J Comp Neurol. 320(4):457-67
- Hornung JP, Fritschy JM (1996) Developmental profile of GABAA-receptors in the marmoset monkey: expression
 of distinct subtypes in pre- and postnatal brain. J Comp Neurol. 367(3):413-30
- Huffman KJ, Krubitzer L. (2001) Thalamo-cortical connections of areas 3a and M1 in marmoset monkeys. J Comp Neurol. 435(3):291-310
- Irle E, Markowitsch HJ (1986) Afferent connections of the substantia innominata/basal nucleus of Meynert in carnivores and primates. J Hirnforsch. 27(3):343-67
- 27. Jouandet ML, Garey LJ, Lipp HP (1984) Distribution of the cells of origin of the corpus callosum and anterior

- commissure in the marmoset monkey. Anat Embryol (Berl). 169(1):45-59
- 28. Kozorovitskiy Y, Hughes M, Lee K, Gould E (2006) Fatherhood affects dendritic spines and vasopressin V1a receptors in the primate prefrontal cortex. Nat Neurosci. 9(9):1094-5
- Maclean CJ, Gaffan D, Baker HF, Ridley RM (2001) Visual discrimination learning impairments produced by combined transections of the anterior temporal stem, amygdala and fornix in marmoset monkeys. Brain Res. 888(1):34-50
- Merkler D, Böscke R, Schmelting B, Czéh B, Fuchs E, Brück W, Stadelmann C (2006) Differential macrophage/ microglia activation in neocortical EAE lesions in the marmoset monkey. Brain Pathol. 16(2):117-23
- Pears A, Parkinson JA, Hopewell L, Everitt BJ, Roberts AC (2003) Lesions of the orbitofrontal but not medial prefrontal cortex disrupt conditioned reinforcement in primates. J Neurosci. 23(35):11189-201
- Pryce CR (2008) Postnatal ontogeny of expression of the corticosteroid receptor genes in mammalian brains: inter-species and intra-species differences. Brain Res Rev. 57(2):596-605
- Ridley RM, Baker JA, Baker HF, Maclean CJ (1994) Restoration of cognitive abilities by cholinergic grafts in cortex of monkeys with lesions of the basal nucleus of Meynert. Neuroscience. 63(3):653-66
- Ridley RM, Baker HF, Leow-Dyke A, Cummings RM (2005) Further analysis of the effects of immunotoxic lesions of the basal nucleus of Meynert reveals substantial impairment on visual discrimination learning in monkeys. Brain Res Bull. 65(5):433-42
- Ridley RM, Barefoot HC, Maclean CJ, Pugh P, Baker HF (1999) Different effects on learning ability after injection of the cholinergic immunotoxin ME20.4lgG-saporin into the diagonal band of Broca, basal nucleus of Meynert, or both in monkeys, Behav Neurosci. 113(2):303-15
- Ridley RM, Maclean CJ, Young FM, Baker HF (2002) Learning impairments in monkeys with combined but not separate excitotoxic lesions of the anterior and mediodorsal thalamic nuclei. Brain Res. 950(1-2):39-51
- Ridley RM, Murray TK, Johnson JA, Baker HF (1986) Learning impairment following lesion of the basal nucleus of Meynert in the marmoset: modification by cholinergic drugs. Brain Res. 376(1):108-16
- Roberts AC, De Salvia MA, Wilkinson LS, Collins P, Muir JL, Everitt BJ, Robbins TW (1994) 6-Hydroxydopamine lesions of the prefrontal cortex in monkeys enhance performance on an analog of the Wisconsin Card Sort Test: possible interactions with subcortical dopamine. J Neurosci. 14(5 Pt 1):2531-44
- Roberts AC, Robbins TW, Everitt BJ, Jones GH, Sirkia TE, Wilkinson J, Page K (1990) The effects of excitotoxic lesions of the basal forebrain on the acquisition, retention and serial reversal of visual discriminations in marmosets, Neuroscience, 34(2):311-29
- Roberts AC, Tomic DL, Parkinson CH, Roeling TA, Cutter DJ, Robbins TW, Everitt BJ (2007) Forebrain connectivity of the prefrontal cortex in the marmoset monkey (Callithrix jacchus): an anterograde and retrograde tract-tracing study. J Comp Neurol. 502(1):86-112
- Roberts AC, Wallis JD (2000) Inhibitory control and affective processing in the prefrontal cortex: neuropsychological studies in the common marmoset. Cereb Cortex. 10(3):252-62
- Rölleke U, Flügge G, Plehm S, Schlumbohm C, Armstrong VW, Dressel R, Uchanska-Ziegler B, Ziegler A, Fuchs E, Czéh B, Walter L (2006) Differential expression of major histocompatibility complex class 1 molecules in the brain of a New World monkey, the common marmoset (Callithrix jacchus). J Neuroimmunol. 176(1-2):39-50
- Schwerdtfeger WK (1979) Direct efferent and afferent connections of the hippocampus with the neocortex in the marmoset monkey. Am J Anat. 156(1):77-82
- 44. Spatz WB (1975) An efferent connection of the solitary cells of Meynert. A study with horseradish peroxidase in the marmoset Callithrix. Brain Res. 92(3):450-5
- Walker SC, Robbins TW, Roberts AC (2008) Differential Contributions of Dopamine and Serotonin to Orbitofrontal Cortex Function in the Marmoset. Cereb Cortex. 19(4):889-98
- Williams SM, Sullivan RK, Scott HL, Finkelstein DI, Colditz PB, Lingwood BE, Dodd PR, Pow DV (2005) Glial glutamate transporter expression patterns in brains from multiple mammalian species. Glia. 49(4):520-41
- Wisniowski L, Ridley RM, Baker HF, Fine A (1992) Tyrosine hydroxylase-immunoreactive neurons in the nucleus basalis of the common marmoset (Callithrix jacchus). J Comp Neurol. 325(3):379-87

Motor cortex

- Burish MJ, Stepniewska I, Kaas JH (2008) Microstimulation and architectonics of frontoparietal cortex in common marmosets (Callithrix jacchus). J Comp Neurol. 507(2):1151-68
- Burman KJ, Palmer SM, Gamberini M, Spitzer MW, Rosa MG (2008) Anatomical and physiological definition of the motor cortex of the marmoset monkey. J Comp Neurol. 506(5):860-76
- Everitt BJ, Sirkiä TE, Roberts AC, Jones GH, Robbins TW (1988) Distribution and some projections of cholinergic neurons in the brain of the common marmoset, Callithrix jacchus. J Comp Neurol. 271(4):533-58
- Fine A, Hoyle C, Maclean CJ, Levatte TL, Baker HF, Ridley RM (1997) Learning impairments following injection of a selective cholinergic immunotoxin, ME20.4 IgG-saporin, into the basal nucleus of Meynert in monkeys. Neuroscience. 81(2):331-43
- Huffman KJ, Krubitzer L (2001) Area 3a: topographic organization and cortical connections in marmoset monkeys. Cereb Cortex. 11(9):849-67
- Huffman KJ, Krubitzer L (2001) Thalamo-cortical connections of areas 3a and M1 in marmoset monkeys. J Comp Neurol. 435(3):291-310

- Hurley MJ, Jackson MJ, Smith LA. Rose S, Jenner P (2005) Immunoautoradiographic analysis of NMDA receptor subunits and associated postsynaptic density proteins in the brain of dyskinetic MPTP-treated common marmosets. Eur J Neurosci. 21(12):3240-50
- Missler M, Wolff JR, Rothe H, Heger W, Merker HJ, Treiber A, Scheid R, Crook GA (1992) Developmental biology of the common marmoset: proposal for a "postnatal staging". J Med Primatol. 21(6):285-98
- Nomoto M, Kaseda S, Iwata S, Shimizu T, Fukuda T, Nakagawa S (2000) The metabolic rate and vulnerability of dopaminergic neurons, and adenosine dynamics in the cerebral cortex, nucleus accumbens, caudate nucleus, and putamen of the common marmoset. J Neurol. 247 Suppl 5:V16-22
- Tsang YM, Chiong F, Kuznetsov D, Kasarskis E. Geula C (2000) Motor neurons are rich in non-phosphorylated neurofilaments: cross-species comparison and alterations in ALS. Brain Res. 861(1):45-58

Somatosensory cortex

- Burish MJ, Stepniewska I, Kaas JH (2008) Microstimulation and architectonics of frontoparietal cortex in common marmosets (Callithrix jacchus). J Comp Neurol. 507(2):1151-68
- Cappe C, Barone P (2005) Heteromodal connections supporting multisensory integration at low levels of cortical processing in the monkey. Eur J Neurosci. 22(11):2886-902
- Coleman GT, Bahramali H, Zhang HQ, Rowe MJ (2001) Characterization of tactile afferent fibers in the hand of the marmoset monkey. J Neurophysiol. 85(5):1793-804
- Coleman GT, Zhang HQ, Murray GM, Zachariah MK, Rowe MJ (1999) Organization of somatosensory areas I and II in marsupial cerebral cortex: parallel processing in the possum sensory cortex. J Neurophysiol. 81(5):2316-24
- Collins CE, Lyon DC, Kaas JH (2005) Distribution across cortical areas of neurons projecting to the superior colliculus in new world monkeys. Anat Rec A Discov Mol Cell Evol Biol. 285(1):619-27
- Fang PC, Jain N, Kaas JH (2002) Few intrinsic connections cross the hand-face border of area 3b of New World monkeys. J Comp Neurol. 454(3):310-9
- Franca JG, Volchan E, Jain N, Catania KC, Oliveira RL, Hess FF, Jablonka M, Rocha-Miranda CE, Kaas JH (2000) Distribution of NADPH-diaphorase cells in visual and somatosensory cortex in four mammalian species. Brain Res. 864(2):163-75
- 8. Hackett TA, Smiley JF, Ulbert I, Karmos G, Lakatos P, de la Mothe LA, Schroeder CE (2007) Sources of somatosensory input to the caudal belt areas of auditory cortex. Perception. 36(10):1419-30
- Huffman KJ, Krubitzer L (2001) Area 3a: topographic organization and cortical connections in marmoset monkeys. Cereb Cortex. 11(9):849-67
- Huffman KJ, Krubitzer L (2001) Thalamo-cortical connections of areas 3a and M1 in marmoset monkeys. J Comp Neurol. 435(3):291-310
- 11. Iyengar S, Qi HX, Jain N, Kaas JH (2007) Cortical and thalamic connections of the representations of the teeth and tongue in somatosensory cortex of new world monkeys. J Comp Neurol. 501(1):95-120
- Kolarik RC, Rasey SK, Wall JT (1994) The consistency, extent, and locations of early-onset changes in cortical nerve dominance aggregates following injury of nerves to primate hands. J Neurosci. 14(7):4269-88
- Krubitzer L, Clarey J, Tweedale R, Elston G, Calford M (1995) A redefinition of somatosensory areas in the lateral sulcus of macaque monkeys. J Neurosci. 15(5 Pt 2):3821-39
- Krubitzer LA, Kaas JH (1988) Responsiveness and somatotopic organization of anterior parietal field 3b and adjoining cortex in newborn and infant monkeys. Somatosens Mot Res. 6(2):179-205
- Krubitzer LA, Kaas JH (1990) The organization and connections of somatosensory cortex in marmosets. J Neurosci. 10(3):952-74
- Krubitzer LA, Kaas JH (1992) The somatosensory thalamus of monkeys: cortical connections and a redefinition of nuclei in marmosets. J Comp Neurol. 319(1):123-40
- Qi HX, Lyon DC, Kaas JH (2002) Cortical and thalamic connections of the parietal ventral somatosensory area in marmoset monkeys (Callithrix jacchus). J Comp Neurol. 443(2):168-82
- Rowe MJ, Turman AB, Murray GM, Zhang HQ (1996) Parallel organization of somatosensory cortical areas I and II for tactile processing. Clin Exp Pharmacol Physiol. 23(10-11):931-8. Review
- Xu J, Wall JT (1996) Cutaneous representations of the hand and other body parts in the cuneate nucleus of a primate, and some relationships to previously described cortical representations. Somatosens Mot Res. 13(3-4):187-97
- Zhang HQ, Murray GM, Coleman GT, Turman AB, Zhang SP, Rowe MJ (2001) Functional characteristics of the parallel SI- and SII-projecting neurons of the thalamic ventral posterior nucleus in the marmoset. J Neurophysiol. 85(5):1805-22
- Zhang HQ, Murray GM, Turman AB, Mackie PD, Coleman GT, Rowe MJ (1996) Parallel processing in cerebral cortex of the marmoset monkey: effect of reversible SI inactivation on tactile responses in SII. J Neurophysiol. 76(6):3633-55
- 22. Zhang HQ, Zachariah MK, Coleman GT, Rowe MJ (2001) Hierarchical equivalence of somatosensory areas I and II for tactile processing in the cerebral cortex of the marmoset monkey. J Neurophysiol. 85(5):1823-35

Visual cortex

- Baldauf ZB (2005) SMI-32 parcellates the visual cortical areas of the marmoset. Neurosci Lett. 383(1-2):109-14
- Barefoot HC, Baker HF, Ridley RM (2000) Synergistic effects of unilateral immunolesions of the cholinergic projections from the basal forebrain and contralateral ablations of the inferotemporal cortex and hippocampus in monkeys. Neuroscience. 98(2):243-51
- Barefoot HC, Maclean CJ, Baker HF, Ridley RM (2003) Unilateral hippocampal and inferotemporal cortex lesions in opposite hemispheres impair learning of single-pair visual discriminations as well as visuovisual conditional tasks in monkeys. Behav Brain Res. 141(1):51-62
- Bourne JA, Lui L, Tweedale R, Rosa MG (2004) First- and second-order stimulus length selectivity in New World monkey striate cortex. Eur J Neurosci. 19(1):169-80
- Bourne JA, Rosa MG (2003) Neurofilament protein expression in the geniculostriate pathway of a New World monkey (Callithrix jacchus). Exp Brain Res. 150(1):19-24
- Bourne JA, Rosa MG (2003) Preparation for the in vivo recording of neuronal responses in the visual cortex of anaesthetised marmosets (Callithrix jacchus). Brain Res Brain Res Protoc. 11(3):168-77
- Bourne JA, Rosa MG (2006) Hierarchical development of the primate visual cortex, as revealed by neurofilament immunoreactivity: early maturation of the middle temporal area (MT). Cereb Cortex. 16(3):405-14
- Bourne JA, Tweedale R, Rosa MG (2002) Physiological responses of New World monkey V1 neurons to stimuli defined by coherent motion. Cereb Cortex. 12(11):1132-45
- Bourne JA, Warner CE, Rosa MG (2005) Topographic and laminar maturation of striate cortex in early postnatal marmoset monkeys, as revealed by neurofilament immunohistochemistry. Cereb Cortex. 15(6):740-8
- Bourne JA, Warner CE, Upton DJ, Rosa MG (2007) Chemoarchitecture of the middle temporal visual area in the marmoset monkey (Callithrix jacchus): laminar distribution of calcium-binding proteins (calbindin, parvalbumin) and nonphosphorylated neurofilament. J Comp Neurol. 500(5):832-49
- 11. Burman KJ, Lui LL, Rosa MG, Bourne JA (2007) Development of non-phosphorylated neurofilament protein expression in neurones of the New World monkey dorsolateral frontal cortex. Eur J Neurosci. 25(6):1767-79
- Burman KJ, Palmer SM, Gamberini M, Rosa MG (2006) Cytoarchitectonic subdivisions of the dorsolateral frontal cortex of the marmoset monkey (Callithrix jacchus), and their projections to dorsal visual areas. J Comp Neurol. 495(2):149-72
- Chappert-Piquemal C, Fonta C, Malecaze F, Imbert M (2001) Ocular dominance columns in the adult New World Monkey Callithrix jacchus. Vis Neurosci. 18(3):407-12
- Collins CE, Lyon DC, Kaas JH (2003) Responses of neurons in the middle temporal visual area after long-standing lesions of the primary visual cortex in adult new world monkeys. J Neurosci. 23(6):2251-64
- Collins CE, Lyon DC, Kaas JH (2005) Distribution across cortical areas of neurons projecting to the superior colliculus in new world monkeys. Anat Rec A Discov Mol Cell Evol Biol. 285(1):619-27
- Costa MS, Britto LR (1997) Calbindin immunoreactivity delineates the circadian visual centers of the brain of the common marmoset (Callithrix jacchus). Brain Res Bull. 43(4):369-73
- 17. Cusick CG, Gould HJ 3rd, Kaas JH (1984) Interhemispheric connections of visual cortex of owl monkeys (Aotus trivirgatus), marmosets (Callithrix jacchus), and galagos (Galago crassicaudatus). J Comp Neurol. 230(3):311-36
- DeBruyn EJ, Casagrande VA (1981) Demonstration of ocular dominance columns in a New World primate by means of monocular deprivation. Brain Res. 207(2):453-8
- Dick A, Kaske A, Creutzfeldt OD (1991) Topographical and topological organization of the thalamocortical projection to the striate and prestriate cortex in the marmoset (Callithrix jacchus). Exp Brain Res. 84(2):233-53
- Elliott T, Shadbolt NR (1999) A neurotrophic model of the development of the retinogeniculocortical pathway induced by spontaneous retinal waves. J Neurosci. 19(18):7951-70
- Elston GN (2003) Pyramidal cell heterogeneity in the visual cortex of the nocturnal New World owl monkey (Aotus trivirgatus). Neuroscience. 117(1):213-9
- Elston GN, Rosa MG (2006) Ipsilateral corticocortical projections to the primary and middle temporal visual areas
 of the primate cerebral cortex: area-specific variations in the morphology of connectionally identified pyramidal
 cells. Eur J Neurosci. 23(12):3337-45
- Elston GN, Rosa MG, Calford MB (1996) Comparison of dendritic fields of layer III pyramidal neurons in striate and extrastriate visual areas of the marmoset: a Lucifer yellow intracellular injection. Cereb Cortex. 6(6):807-13
- Elston GN, Tweedale R, Rosa MG (1999) Cellular heterogeneity in cerebral cortex: a study of the morphology of pyramidal neurones in visual areas of the marmoset monkey. J Comp Neurol. 415(1):33-51
- Fitzgibbon T, Szmajda BA, Martin PR (2007) First order connections of the visual sector of the thalamic reticular nucleus in marmoset monkeys (Callithrix jacchus). Vis Neurosci. 24(6):857-74
- Fonta C, Chappert C, Imbert M (1997) N-methyl-D-aspartate subunit R1 involvement in the postnatal organization
 of the primary visual cortex of Callithrix jacchus. J Comp Neurol. 386(2):260-76
- Forte JD, Blessing EM, Buzás P, Martin PR (2006) Contribution of chromatic aberrations to color signals in the primate visual system. J Vis. 6(2):97-105
- Forte JD, Hashemi-Nezhad M, Dobbie WJ, Dreher B, Martin PR (2005) Spatial coding and response redundancy in parallel visual pathways of the marmoset Callithrix jacchus. Vis Neurosci. 22(4):479-91
- Franca JG, Volchan E, Jain N, Catania KC, Oliveira RL, Hess FF, Jablonka M, Rocha-Miranda CE, Kaas JH (2000)
 Distribution of NADPH-diaphorase cells in visual and somatosensory cortex in four mammalian species. Brain Res. 864(2):163-75

- Fritsches KA, Rosa MG (1996) Visuotopic organisation of striate cortex in the marmoset monkey (Callithrix jacchus). J Comp Neurol. 372(2):264-82
- Fritschy JM, Garey LJ (1986) Postnatal development of quantitative morphological parameters in the lateral geniculate nucleus of the marmoset monkey. Brain Res. 395(2):157-68
- 32. Fritschy JM, Garey LJ (1986) Quantitative changes in morphological parameters in the developing visual cortex of the marmoset monkey. Brain Res. 394(2):173-88
- 33. Garey LJ, de Courten C (1983) Structural development of the lateral geniculate nucleus and visual cortex in monkey and man, Behav Brain Res. 10(1):3-13
- 34. Gebhard R, Zilles K, Schleicher A, Everitt BJ, Robbins TW, Divac I (1993) Distribution of seven major neurotransmitter receptors in the striate cortex of the New World monkey Callithrix jacchus. Neuroscience. 56(4):877-85
- 35. Goodchild AK, Martin PR (1998) The distribution of calcium-binding proteins in the lateral geniculate nucleus and visual cortex of a New World monkey, the marmoset, Callithrix jacchus. Vis Neurosci. 15(4):625-42
- Guo K, Robertson R, Nevado A, Pulgarin M, Mahmoodi S, Young MP (2006) Primary visual cortex neurons that contribute to resolve the aperture problem. Neuroscience. 138(4):1397-406
- Jeffs J, Ichida JM, Federer F, Angelucci A (2008) Anatomical Evidence for Classical and Extra-classical Receptive Field Completion Across the Discontinuous Horizontal Meridian Representation of Primate Area V2. Cereb Cortex. 16
- Kaske A, Dick A, Creutzfeldt OD (1991) The local domain for divergence of subcortical afferents to the striate and extrastriate visual cortex in the common marmoset (Callithrix jacchus): a multiple labelling study. Exp Brain Res. 84(2):254-65
- Kilavik BE, Silveira LC, Kremers J (2007) Spatial receptive field properties of lateral geniculate cells in the owl monkey (Aotus azarae) at different contrasts: a comparative study. Eur J Neurosci. 26(4):992-1006
- Kunz B, Spatz WB (1985) A callosal projection of area 17 upon the border region of area MT in the marmoset monkey, Callithrix jacchus. J Comp Neurol. 239(4):413-9
- Laffont-Proust I, Fonta C, Renaud L, Hässig R, Moya KL (2007) Developmental changes in cellular prion protein in primate visual cortex. J Comp Neurol. 504(6):646-58
- Liu GB, Pettigrew JD (2003) Orientation mosaic in barn owl's visual Wulst revealed by optical imaging: comparison with cat and monkey striate and extra-striate areas. Brain Res. 961(1):153-8
- Lui LL, Bourne JA, Rosa MG (2005) Single-unit responses to kinetic stimuli in New World monkey area V2: physiological characteristics of cue-invariant neurones. Exp Brain Res. 162(1):100-8
- Lui LL, Bourne JA, Rosa MG (2006) Functional response properties of neurons in the dorsomedial visual area of New World monkeys (Callithrix jacchus). Cereb Cortex. 16(2):162-77
- 45. Lui LL, Bourne JA, Rosa MG (2007) Spatial and temporal frequency selectivity of neurons in the middle temporal visual area of new world monkeys (Callithrix jacchus). Eur J Neurosci. 25(6):1780-92
- Lui LL, Bourne JA, Rosa MG (2007) Spatial summation, end inhibition and side inhibition in the middle temporal visual area (MT). J Neurophysiol. 97(2):1135-48
- Lyon DC, Kaas JH (2001) Connectional and architectonic evidence for dorsal and ventral V3, and dorsomedial area in marmoset monkeys. J Neurosci. 21(1):249-61
- 48. Lyon DC, Kaas JH (2002) Connectional evidence for dorsal and ventral V3, and other extrastriate areas in the prosimian primate, Galago garnetti. Brain Behav Evol. 59(3):114-29
- Malach R (1992) Dendritic sampling across processing streams in monkey striate cortex. J Comp Neurol. 315(3):303-12
- Markstahler U, Bach M, Spatz WB (1998) Transient molecular visualization of ocular dominance columns (ODCs) in normal adult marmosets despite the desegregated termination of the retino-geniculo-cortical pathways. J Comp Neurol, 393(1):118-34
- Martin PR, White AJ, Goodchild AK, Wilder HD, Sefton AE (1997) Evidence that blue-on cells are part of the third geniculocortical pathway in primates. Eur J Neurosci. 9(7):1536-41
- McLoughlin N, Schiessl I (2006) Orientation selectivity in the common marmoset (Callithrix jacchus): the periodicity of orientation columns in V1 and V2, Neuroimage. 31(1):76-85
- Missler M, Eins S, Böttcher H, Wolff JR (1994) Postnatal development of glial fibrillary acidic protein, vimentin and S100 protein in monkey visual cortex: evidence for a transient reduction of GFAP immunoreactivity. Brain Res Dev Brain Res. 82(1-2):103-17
- Missler M, Eins S, Merker HJ, Rothe H, Wolff JR (1993) Pre- and postnatal development of the primary visual cortex of the common marmoset. I. A changing space for synaptogenesis. J Comp Neurol. 333(1):41-52
- Missler M, Wolff A, Merker HJ, Wolff JR (1993) Pre- and postnatal development of the primary visual cortex of the common marmoset. II. Formation, remodelling, and elimination of synapses as overlapping processes. J Comp Neurol. 333(1):53-67
- Palmer SM, Rosa MG (2006) A distinct anatomical network of cortical areas for analysis of motion in far peripheral vision. Eur J Neurosci. 24(8):2389-405
- Palmer SM, Rosa MG (2006) Quantitative analysis of the corticocortical projections to the middle temporal area in the marmoset monkey: evolutionary and functional implications. Cereb Cortex. 16(9):1361-75
- Ridley RM, Baker HF, Cummings RM, Green ME, Leow-Dyke A (2005) Mild topographical memory impairment following crossed unilateral lesions of the mediodorsal thalamic nucleus and the inferotemporal cortex. Behav Neurosci. 119(2):518-25

- Ridley RM, Baker HF, Leow-Dyke A. Cummings RM (2005) Further analysis of the effects of immunotoxic lesions of the basal nucleus of Meynert reveals substantial impairment on visual discrimination learning in monkeys. Brain Res Bull. 65(5):433-42
- Ridley RM, Baker HF, Mills DA, Green ME, Cummings RM (2004) Topographical memory impairments after unilateral lesions of the anterior thalamus and contralateral inferotemporal cortex. Neuropsychologia. 42(9):1178-91
- Ridley RM, Warner KA, Maclean CJ, Gaffan D, Baker HF (2001) Visual agnosia and Klüver-Bucy syndrome in marmosets (Callithrix jacchus) following ablation of inferotemporal cortex, with additional mnemonic effects of immunotoxic lesions of cholinergic projections to medial temporal areas. Brain Res. 898(1):136-51
- Roberts AC, Robbins TW, Everitt BJ, Jones GH, Sirkia TE, Wilkinson J, Page K (1990) The effects of excitotoxic lesions of the basal forebrain on the acquisition, retention and serial reversal of visual discriminations in marmosets. Neuroscience. 34(2):311-29
- 63. Roberts MJ, Zinke W, Guo K, Robertson R, McDonald JS, Thiele A (2005) Acetylcholine dynamically controls spatial integration in marmoset primary visual cortex. J Neurophysiol. 93(4):2062-72
- Roe AW, Fritsches K, Pettigrew JD (2005) Optical imaging of functional organization of V1 and V2 in marmoset visual cortex. Anat Rec A Discov Mol Cell Evol Biol. 287(2):1213-25
- Roe AW, Fritsches K, Pettigrew JD (2005) Optical imaging of functional organization of V1 and V2 in marmoset visual cortex. Anat Rec A Discov Mol Cell Evol Biol. 287(2):1213-25
- 66. Roe AW, Fritsches K, Pettigrew JD (2005) Optical imaging of functional organization of V1 and V2 in marmoset visual cortex. Anat Rec A Discov Mol Cell Evol Biol. 287(2):1213-25
- Rosa MG, Elston GN (1998) Visuotopic organisation and neuronal response selectivity for direction of motion in visual areas of the caudal temporal lobe of the marmoset monkey (Callithrix jacchus): middle temporal area, middle temporal crescent, and surrounding cortex. J Comp Neurol. 393(4):505-27
- Rosa MG, Fritsches KA, Elston GN (1997) The second visual area in the marmoset monkey: visuotopic organisation, magnification factors, architectonical boundaries, and modularity. J Comp Neurol. 387(4):547-67
- Rosa MG, Palmer SM, Gamberini M, Tweedale R, Piñon MC, Bourne JA (2005) Resolving the organization of the New World monkey third visual complex: the dorsal extrastriate cortex of the marmoset (Callithrix jacchus). J Comp Neurol, 483(2):164-91
- Rosa MG, Schmid LM (1995) Visual areas in the dorsal and medial extrastriate cortices of the marmoset. J Comp Neurol. 359(2):272-99
- Rosa MG, Tweedale R (2000) Visual areas in lateral and ventral extrastriate cortices of the marmoset monkey. J Comp Neurol. 422(4):621-51
- Rosa MG, Tweedale R. Elston GN (2000) Visual responses of neurons in the middle temporal area of new world monkeys after lesions of striate cortex. J Neurosci. 20(14):5552-63
- Schiessl I, McLoughlin N (2003) Optical imaging of the retinotopic organization of V1 in the common marmoset. Neuroimage. 20(3):1857-64
- Sengpiel F, Troilo D, Kind PC, Graham B, Blakemore C (1996) Functional architecture of area 17 in normal and monocularly deprived marmosets (Callithrix jacchus). Vis Neurosci. 13(1):145-60
- 75. Solomon SG (2002) Striate cortex in dichromatic and trichromatic marmosets: neurochemical compartmentalization and geniculate input. J Comp Neurol. 450(4):366-81
- 76. Spatz WB (1975) Thalamic and other subcortical projections to area MT (visual area of superior temporal sulcus) in the marmoset Callithrix jacchus, Brain Res. 99(1):129-34
- Spatz WB (1977) Topographically organized reciprocal connections between areas 17 and MT (visual area of superior temporal sulcus) in the marmoset Callithrix jacchus. Exp Brain Res. 27(5):559-72
- Spatz WB (1978) The retino-geniculo-cortical pathway in Callithrix. I. Intraspecific variations in the lamination pattern of the lateral geniculate nucleus. Exp Brain Res. 33(3-4):551-63
- Spatz WB (1979) The retino-geniculo-cortical pathway in Callithrix. II. The geniculo-cortical projection. Exp Brain Res. 36(3):401-10
- Spatz WB (1989) Loss of ocular dominance columns with maturity in the monkey, Callithrix jacchus. Brain Res. 488(1-2):376-80
- 81. Spatz WB, Erdmann G (1974) Striate cortex projections to the lateral geniculate and other thalamic nuclei: a study using degeneration and autoradiographic tracing methods in the marmoset Callithrix. Brain Res. 82(1):91-108
- 82. Spatz WB, Illing RB, Weisenhorn DM (1994) Distribution of cytochrome oxidase and parvalbumin in the primary visual cortex of the adult and neonate monkey, Callithrix jacchus. J Comp Neurol. 339(4):519-34
- Spatz WB, Kunz B (1984) Area 17 of anthropoid primates does participate in visual callosal connections, Neurosci Lett. 48(1):49-53
- Spatz WB, Kunz B, Steffen H (1987) A new heterotopic callosal projection of primary visual cortex in the monkey, Callithrix jacchus. Brain Res. 403(1):158-61
- Spatz WB, Tigges J (1972) Experimental-anatomical studies on the "middle temporal visual area (MT)" in primates.
 Efferent cortico-cortical connections in the marmoset Callithrix jacchus. J Comp Neurol. 146(4):451-64
- Spatz WB, Vogt DM, Illing RB (1993) Distribution of cytochrome oxidase in layers IV and V of the striate cortex in neonate monkeys. Exp Brain Res. 95(1):183-6
- Stepniewska I, Ql HX, Kaas JH (2000) Projections of the superior colliculus to subdivisions of the inferior pulvinar in New World and Old World monkeys. Vis Neurosci. 17(4):529-49
- 88. Stevens CF (2002) Predicting functional properties of visual cortex from an evolutionary scaling law. Neuron.

- 36(1):139-42
- 89. Szmajda BA, Grünert U, Martin PR (2008) Retinal ganglion cell inputs to the koniocellular pathway. J Comp Neurol. 510(3):251-68
- Takahata T, Komatsu Y, Watakabe A, Hashikawa T, Tochitani S. Yamamori T (2006) Activity-dependent expression of occ1 in excitatory neurons is a characteristic feature of the primate visual cortex. Cereb Cortex. 16(7):929-40
- Tinsley CJ, Webb BS, Barraclough NE, Vincent CJ, Parker A, Derrington AM (2003) The nature of V1 neural responses to 2D moving patterns depends on receptive-field structure in the marmoset monkey. J Neurophysiol. 90(2):930-7
- vogt Weisenhorn DM, Illing RB, Spatz WB (1995) Morphology and connections of neurons in area 17 projecting to the extrastriate areas MT and 19DM and to the superior colliculus in the monkey Callithrix jacchus. J Comp Neurol. 362(2):233-55
- Webb BS, Tinsley CJ, Barraclough NE, Easton A, Parker A, Derrington AM (2002) Feedback from V1 and inhibition from beyond the classical receptive field modulates the responses of neurons in the primate lateral geniculate nucleus. Vis Neurosci. 19(5):583-92
- Zinke W, Roberts MJ, Guo K, McDonald JS, Robertson R, Thiele A (2006) Cholinergic modulation of response properties and orientation tuning of neurons in primary visual cortex of anaesthetized Marmoset monkeys. Eur J Neurosci. 24(1):314-28

Auditory cortex

- Aitkin LM, Kudo M, Irvine DR (1988) Connections of the primary auditory cortex in the common marmoset, Callithrix jacchus. J Comp Neurol. 269(2):235-48
- de la Mothe LA, Blumell S, Kajikawa Y, Hackett TA (2006) Cortical connections of the auditory cortex in marmoset monkeys: core and medial belt regions. J Comp Neurol. 496(1):27-71
- Eliades SJ, Wang X (2005) Dynamics of auditory-vocal interaction in monkey auditory cortex. Cereb Cortex. 15(10):1510-23
- Hackett TA, Smiley JF, Ulbert I, Karmos G, Lakatos P, de la Mothe LA, Schroeder CE (2007) Sources of somatosensory input to the caudal belt areas of auditory cortex. Perception. 36(10):1419-30
- Walker KM, Ahmed B, Schnupp JW (2008) Linking cortical spike pattern codes to auditory perception. J Cogn Neurosci. 20(1):135-52

Association cortex

- Elston GN, Rosa MG (2006) Ipsilateral corticocortical projections to the primary and middle temporal visual areas of the primate cerebral cortex: area-specific variations in the morphology of connectionally identified pyramidal cells. Eur J Neurosci. 23(12):3337-45
- 2. Palmer SM, Rosa MG (2006) Quantitative analysis of the corticocortical projections to the middle temporal area in the marmoset monkey; evolutionary and functional implications. Cereb Cortex. 16(9):1361-75
- 3. Spatz WB (1977) Topographically organized reciprocal connections between areas 17 and MT (visual area of superior temporal sulcus) in the marmoset Callithrix jacchus. Exp Brain Res. 27(5):559-72
- Zhang HQ, Murray GM, Turman AB, Mackie PD, Coleman GT. Rowe MJ (1996) Parallel processing in cerebral cortex of the marmoset monkey: effect of reversible SI inactivation on tactile responses in SII. J Neurophysiol. 76(6):3633-55
- 5. Zhang HQ, Zachariah MK, Coleman GT, Rowe MJ (2001) Hierarchical equivalence of somatosensory areas 1 and II for tactile processing in the cerebral cortex of the marmoset monkey. J Neurophysiol. 85(5):1823-35

Hippocampus

- Abrahám H, Czéh B, Fuchs E, Seress L (2005) Mossy cells and different subpopulations of pyramidal neurons are immunoreactive for cocaine- and amphetamine-regulated transcript peptide in the hippocampal formation of nonhuman primates and tree shrew (Tupaia belangeri). Neuroscience. 136(1):231-40
- Barefoot HC, Maclean CJ, Baker HF, Ridley RM (2003) Unilateral hippocampal and inferotemporal cortex lesions in opposite hemispheres impair learning of single-pair visual discriminations as well as visuovisual conditional tasks in monkeys. Behav Brain Res. 141(1):51-62
- 3. Barinaga M (1998) No-new-neurons dogma loses ground. Science. 279(5359):2041-2
- Bittencourt S, Pereira CM, Avedissian M, Delamano A, Mello LE, Castilho BA (2008) Distribution of the protein IMPACT, an inhibitor of GCN2, in the mouse, rat, and marmoset brain. J Comp Neurol. 507(5):1811-30
- Brown AW, Verschoyle RD, Street BW, Aldridge WN, Grindley H (1984) The neurotoxicity of trimethyltin chloride in hamsters, gerbils and marmosets. J Appl Toxicol. 4(1):12-21
- 6. Del Bigio MR (1999) Proliferative status of cells in adult human dentate gyrus. Microsc Res Tech. 45(6):353-8
- 7. Del Zompo M, Piccardi MP, Ruiu S, Albanese A, Morelli M (1992) Localization of MPP+ binding sites in the brain of various mammalian species. J Neural Transm Park Dis Dement Sect. 4(3):181-90
- 8. Dumont Y, Jacques D, Bouchard P, Quirion R (1998) Species differences in the expression and distribution of the

- neuropeptide Y Y1, Y2, Y4, and Y5 receptors in rodents, guinea pig, and primates brains, J Comp Neurol, 402(3):372-84
- Dwyer BE, Fujikawa DG, Wasterlain CG (1986) Metabolic anatomy of generalized bicuculline seizures in the newborn marmoset monkey. Exp Neurol. 94(1):213-27
- Fujikawa DG, Söderfeldt B, Wasterlain CG (1992) Neuropathological changes during generalized seizures in newborn monkeys. Epilepsy Res. 12(3):243-51
- Gould E, Reeves AJ, Fallah M, Tanapat P, Gross CG, Fuchs E (1999) Hippocampal neurogenesis in adult Old World primates. Proc Natl Acad Sci U S A. 96(9):5263-7
- Gould E, Tanapat P, McEwen BS. Flügge G, Fuchs E (1998) Proliferation of granule cell precursors in the dentate gyrus of adult monkeys is diminished by stress. Proc Natl Acad Sci U S A. 95(6):3168-71
- Gozzo S, Perretta G (1992) The mossy fiber terminals in the hippocampal formation of Callithrix jacchus: comparative and evolutionary considerations. Int J Neurosci. 63(3-4):163-77
- Gozzo S, Perretta G, Monaco V, Andreozzi U, Rossiello E (1993) The neuropathology of trimethyltin in the marmoset (Callithrix jacchus) hippocampal formation. Ecotoxicol Environ Saf. 26(3):293-301
- Hornung JP, Celio MR (1992) The selective innervation by serotoninergic axons of calbindin-containing interneurons in the neocortex and hippocampus of the marmoset. J Comp Neurol. 320(4):457-67
- Hurley MJ, Jackson MJ, Smith LA, Rose S, Jenner P (2005) Immunoautoradiographic analysis of NMDA receptor subunits and associated postsynaptic density proteins in the brain of dyskinetic MPTP-treated common marmosets. Eur J Neurosci. 21(12):3240-50
- Johnson EO, Brady L, Gold PW, Chrousos GP (1996) Distribution of hippocampal mineralocorticoid and glucocorticoid receptor mRNA in a glucocorticoid resistant nonhuman primate. Steroids. 61(2):69-73
- 18. Kempermann G, Gage FH (1998) Closer to neurogenesis in adult humans. Nat Med. 4(5):555-7
- Korr H (1976) Autoradiographic studies on the distribution of 3H-mescaline in the brain of the marmoset, Callithrix jacchus. Psychopharmacologia. 46(1):115-7
- Kozorovitskiy Y, Gross CG, Kopil C, Battaglia L, McBreen M, Stranahan AM, Gould E (2005) Experience induces structural and biochemical changes in the adult primate brain. Proc Natl Acad Sci U S A, 102(48):17478-82
- Kraemer M, Zilles K, Schleicher A, Gebhard R, Robbins TW, Everitt BJ, Divac 1 (1995) Quantitative receptor autoradiography of eight different transmitter-binding sites in the hippocampus of the common marmoset, Callithrix jacchus. Anat Embryol (Berl). 191(3):213-25
- Leuner B, Kozorovitskiy Y, Gross CG, Gould E (2007) Diminished adult neurogenesis in the marmoset brain precedes old age. Proc Natl Acad Sci U S A, 104(43):17169-73
- Maclean CJ, Baker HF, Fine A, Ridley RM (1997) The distribution of p75 neurotrophin receptor-immunoreactive cells in the forebrain of the common marmoset (Callithrix jacchus). Brain Res Bull. 43(2):197-208
- Pothuizen HH, Feldon J, Jongen-Rêlo AL (2004) Co-expression of calretinin and gamma-aminobutyric acid in neurons of the entorhinal cortex of the common marmoset monkey. Hippocampus. 14(5):615-27
- Pryce CR (2008) Postnatal ontogeny of expression of the corticosteroid receptor genes in mammalian brains: interspecies and intra-species differences. Brain Res Rev. 57(2):596-605
- Pryce CR, Feldon J, Fuchs E, Knuesel I, Oertle T, Sengstag C, Spengler M, Weber E, Weston A, Jongen-Rélo A (2005) Postnatal ontogeny of hippocampal expression of the mineralocorticoid and glucocorticoid receptors in the common marmoset monkey. Eur J Neurosci. 21(6):1521-35
- Ridley RM, Baker HF, Harder JA. Pearson C (1996) Effects of lesions of different parts of the septo-hippocampal system in primates on learning and retention of information acquired before or after surgery. Brain Res Bull. 40(1):21-32
- Ridley RM, Baker HF, Leow-Dyke A, Cummings RM (2005) Further analysis of the effects of immunotoxic lesions
 of the basal nucleus of Meynert reveals substantial impairment on visual discrimination learning in monkeys. Brain
 Res Bull. 65(5):433-42
- Ridley RM, Baker HF, Mills DA, Green ME, Cummings RM (2004) Topographical memory impairments after unilateral lesions of the anterior thalamus and contralateral inferotemporal cortex. Neuropsychologia. 42(9):1178-91
- Ridley RM, Barefoot HC, Maclean CJ, Pugh P, Baker HF (1999) Different effects on learning ability after injection of the cholinergic immunotoxin ME20.4IgG-saporin into the diagonal band of Broca, basal nucleus of Meynert, or both in monkeys. Behav Neurosci. 113(2):303-15
- Ridley RM, Maclean CJ, Young FM, Baker HF (2002) Learning impairments in monkeys with combined but not separate excitotoxic lesions of the anterior and mediodorsal thalamic nuclei. Brain Res. 950(1-2):39-51
- Ridley RM, Timothy CJ, Maclean CJ, Baker HF (1995) Conditional learning and memory impairments following neurotoxic lesion of the CA1 field of the hippocampus. Neuroscience. 67(2):263-75
- Ridley RM, Warner KA, Maclean CJ, Gaffan D, Baker HF (2001) Visual agnosia and Klüver-Bucy syndrome in marmosets (Callithrix jacchus) following ablation of inferotemporal cortex, with additional mnemonic effects of immunotoxic lesions of cholinergic projections to medial temporal areas. Brain Res. 898(1):136-51
- Schwerdtfeger WK (1979) Direct efferent and afferent connections of the hippocampus with the neocortex in the marmoset monkey. Am J Anat. 156(1):77-82
- Schwerdtfeger WK (1984) Structure and fiber connections of the hippocampus. A comparative study. Adv Anat Embryol Cell Biol. 83:1-74
- Seress L, Abrahám H, Czéh B, Fuchs E, Léránth C (2008) Calretinin expression in hilar mossy cells of the hippocampal dentate gyrus of nonhuman primates and humans. Hippocampus. 18(4):425-34

- 37. Tauber SC, Schlumbohm C, Schilg L, Fuchs E, Nau R, Gerber J (2006) Intrauterine exposure to dexamethasone impairs proliferation but not neuronal differentiation in the dentate gyrus of newborn common marmoset monkeys. Brain Pathol. 16(3):209-17
- West MJ, Schwerdtfeger WK (1985) An allometric study of hippocampal components. A comparative study of the brains of the European hedgehog (Erinaceus europaeus), the tree shrew (Tupaia glis), and the marmoset monkey (Callithrix jacchus). Brain Behav Evol. 27(2-4):93-105
- Yau JL, Rasmuson S, Andrew R, Graham M, Noble J, Olsson T, Fuchs E, Lathe R, Seckl JR (2003) Dehydroepiandrosterone 7-hydroxylase CYP7B: predominant expression in primate hippocampus and reduced expression in Alzheimer's disease. Neuroscience. 121(2):307-14

Amygdala

- Braesicke K, Parkinson JA, Reekie Y, Man MS, Hopewell L, Pears A, Crofts H, Schnell CR, Roberts AC (2005) Autonomic arousal in an appetitive context in primates: a behavioural and neural analysis. Eur J Neurosci. 21(6):1733-40
- Brevard ME, Meyer JS, Harder JA, Ferris CF (2006) Imaging brain activity in conscious monkeys following oral MDMA ("ecstasy"). Magn Reson Imaging. 24(6):707-14
- Costall B, Naylor RJ, Tyers MB (1990) The psychopharmacology of 5-HT3 receptors. Pharmacol Ther. 47(2):181-202
- 4. Irle E, Markowitsch HJ (1986) Afferent connections of the substantia innominata/basal nucleus of Meynert in carnivores and primates. J Hirnforsch. 27(3):343-67
- 5. Maclean CJ, Baker HF, Fine A, Ridley RM (1997) The distribution of p75 neurotrophin receptor-immunoreactive cells in the forebrain of the common marmoset (Callithrix jacchus). Brain Res Bull. 43(2):197-208
- Maclean CJ, Gaffan D, Baker HF, Ridley RM (2001) Visual discrimination learning impairments produced by combined transections of the anterior temporal stem, amygdala and fornix in marmoset monkeys. Brain Res. 888(1):34-50
- Markowitsch HJ, Emmans D, Irle E, Streicher M, Preilowski B (1985) Cortical and subcortical afferent connections of the primate's temporal pole: a study of rhesus monkeys, squirrel monkeys, and marmosets. J Comp Neurol. 242(3):425-58
- Meyer JS, Brevard ME, Piper BJ, Ali SF, Ferris CF (2006) Neural effects of MDMA as determined by functional magnetic resonance imaging and magnetic resonance spectroscopy in awake marmoset monkeys. Ann N Y Acad Sci. 1074:365-76
- Parkinson JA, Crofts HS, McGuigan M, Tomic DL, Everitt BJ, Roberts AC (2001) The role of the primate amygdala in conditioned reinforcement. J Neurosci. 21(19):7770-80
- Pears A, Parkinson JA, Hopewell L, Everitt BJ, Roberts AC (2003) Lesions of the orbitofrontal but not medial prefrontal cortex disrupt conditioned reinforcement in primates. J Neurosci. 23(35):11189-201
- Pryce CR (2008) Postnatal ontogeny of expression of the corticosteroid receptor genes in mammalian brains: interspecies and intra-species differences. Brain Res Rev. 57(2):596-605
- 12. Ridley RM, Murray TK, Johnson JA, Baker HF (1986) Learning impairment following lesion of the basal nucleus of Meynert in the marmoset: modification by cholinergic drugs. Brain Res. 376(1):108-16
- Roberts AC, Robbins TW, Everitt BJ, Jones GH, Sirkia TE, Wilkinson J, Page K (1990) The effects of excitotoxic lesions of the basal forebrain on the acquisition, retention and serial reversal of visual discriminations in marmosets. Neuroscience. 34(2):311-29
- Roberts AC, Robbins TW, Everitt BJ, Muir JL (1992) A specific form of cognitive rigidity following excitotoxic lesions of the basal forebrain in marmosets. Neuroscience. 47(2):251-64
- Roberts AC, Tomic DL, Parkinson CH, Roeling TA, Cutter DJ, Robbins TW, Everitt BJ (2007) Forebrain connectivity of the prefrontal cortex in the marmoset monkey (Callithrix jacchus): an anterograde and retrograde tract-tracing study. J Comp Neurol. 502(1):86-112
- Wang Z, Moody K, Newman JD, Insel TR (1997) Vasopressin and oxytocin immunoreactive neurons and fibers in the forebrain of male and female common marmosets (Callithrix jacchus). Synapse. 27(1):14-25
- Wang Z, Toloczko D, Young LJ, Moody K, Newman JD, Insel TR (1997) Vasopressin in the forebrain of common marmosets (Callithrix jacchus): studies with in situ hybridization, immunocytochemistry and receptor autoradiography. Brain Res. 768(1-2):147-56

Diencephalon

- Aitkin L, Park V (1993) Audition and the auditory pathway of a vocal New World primate, the common marmoset. Prog Neurobiol. 41(3):345-67. Review
- 2. Bartlett EL, Wang X (2007) Neural representations of temporally modulated signals in the auditory thalamus of awake primates. J Neurophysiol. 97(2):1005-17
- Blessing EM, Solomon SG, Hashemi-Nezhad M, Morris BJ, Martin PR (2004) Chromatic and spatial properties of parvocellular cells in the lateral geniculate nucleus of the marmoset (Callithrix jacchus). J Physiol. 557(Pt 1):229-45
- 4. Bourne JA, Rosa MG (2003) Neurofilament protein expression in the geniculostriate pathway of a New World

- monkey (Callithrix jacchus). Exp Brain Res. 150(1):19-24
- Brysch I, Brysch W, Creutzfeldt O, Hayes NL, Schlingensiepen KH (1984) The second, intralaminar thalamocortical projection system. Anat Embryol (Berl). 169(2):111-8
- Brysch W, Brysch I, Creutzfeldt OD, Schlingensiepen R, Schlingensiepen KH (1990) The topology of the thalamocortical projections in the marmoset monkey (Callithrix jacchus). Exp Brain Res. 81(1):1-17
- Cavalcante JS, Alves AS, Costa MS, Britto LR (2002) Differential distribution of afferents containing serotonin and neuropeptide Y within the marmoset suprachiasmatic nucleus. Brain Res. 927(2):200-3
- Cavalcante JS, Britto LR, Toledo CA, Nascimento ES Jr, Lima RR, Pontes AL, Costa MS (2008) Calcium-binding proteins in the circadian centers of the common marmoset (Callithrix jacchus) and the rock cavy (Kerodon rupestris) brains. Brain Res Bull. 76(4):354-60
- Cavalcante JS, Costa MS, Santee UR, Britto LR (2005) Retinal projections to the midline and intralaminar thalamic nuclei in the common marmoset (Callithrix jacchus). Brain Res. 1043(1-2):42-7
- Chappert-Piquemal C, Fonta C, Malecaze F, Imbert M (2001) Ocular dominance columns in the adult New World Monkey Callithrix jacchus. Vis Neurosci. 18(3):407-12
- Chevassus-au-Louis N, Cooper HM (1998) Is there a geniculohypothalamic tract in primates? A comparative immunohistochemical study in the circadian system of strepsirhine and haplorhine species. Brain Res. 805(1-2):213-9
- Costa MS, Britto LR (1997) Calbindin immunoreactivity delineates the circadian visual centers of the brain of the common marmoset (Callithrix jacchus). Brain Res Bull. 43(4):369-73
- Costa MS, Santee UR, Cavalcante JS, Moraes PR, Santos NP, Britto LR (1999) Retinohypothalamic projections in the common marmoset (Callithrix jacchus): A study using cholera toxin subunit B. J Comp Neurol. 415(3):393-403
- de la Mothe LA, Blumell S, Kajikawa Y, Hackett TA (2006) Thalamic connections of the auditory cortex in marmoset monkeys: core and medial belt regions. J Comp Neurol. 496(1):72-96
- DeBruyn EJ, Casagrande VA (1981) Demonstration of ocular dominance columns in a New World primate by means of monocular deprivation. Brain Res. 207(2):453-8
- Dick A, Kaske A, Creutzfeldt OD (1991) Topographical and topological organization of the thalamocortical projection to the striate and prestriate cortex in the marmoset (Callithrix jacchus). Exp Brain Res. 84(2):233-53
- Elliott T, Shadbolt NR (1999) A neurotrophic model of the development of the retinogeniculocortical pathway induced by spontaneous retinal waves. J Neurosci. 19(18):7951-70
- Felisberti F, Derrington AM (2001) Long-range interactions in the lateral geniculate nucleus of the New-World monkey, Callithrix jacchus. Vis Neurosci. 18(2):209-18
- FitzGibbon T, Solomon SG, Goodchild AK (2000) Distribution of calbindin, parvalbumin, and calretinin immunoreactivity in the reticular thalamic nucleus of the marmoset: evidence for a medial leaflet of incertal neurons. Exp Neurol. 164(2):371-83
- Fitzgibbon T, Szmajda BA, Martin PR (2007) First order connections of the visual sector of the thalamic reticular nucleus in marmoset monkeys (Callithrix jacchus). Vis Neurosci. 24(6):857-74
- Fonta C, Chappert C, Imbert M (1997) N-methyl-D-aspartate subunit R1 involvement in the postnatal organization
 of the primary visual cortex of Callithrix jacchus. J Comp Neurol. 386(2):260-76
- Fonta C, Chappert C, Imbert M (2000) Effect of monocular deprivation on NMDAR1 immunostaining in ocular dominance columns of the marmoset Callithrix jacchus. Vis Neurosci. 17(3):345-52
- 23. Fritschy JM, Garey LJ (1986) Postnatal development of quantitative morphological parameters in the lateral geniculate nucleus of the marmoset monkey. Brain Res. 395(2):157-68
- Fritschy JM, Garey LJ (1988) Postnatal development of dendrites of relay neurons in the lateral geniculate nucleus of the marmoset (Callithrix jacchus): a quantitative Golgi study. J Comp Neurol. 268(2):234-47
- Garey LJ, de Courten C (1983) Structural development of the lateral geniculate nucleus and visual cortex in monkey and man. Behav Brain Res. 10(1):3-13
- Gibb WR, Lees AJ, Jenner P, Marsden CD (1986) The dopamine neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) produces histological lesions in the hypothalamus of the common marmoset. Neurosci Lett. 65(1):79-83
- 27. Goodchild AK, Martin PR (1998) The distribution of calcium-binding proteins in the lateral geniculate nucleus and visual cortex of a New World monkey, the marmoset, Callithrix jacchus. Vis Neurosci. 15(4):625-42
- Gouardères C, Puget A, Zajac JM (2004) Detailed distribution of neuropeptide FF receptors (NPFF1 and NPFF2) in the rat, mouse, octodon, rabbit, guinea pig, and marmoset monkey brains: a comparative autoradiographic study. Synapse. 51(4):249-69
- Hardman CD, Henderson JM, Finkelstein DI, Horne MK, Paxinos G, Halliday GM (2002) Comparison of the basal ganglia in rats, marmosets, macaques, baboons, and humans: volume and neuronal number for the output, internal relay, and striatal modulating nuclei. J Comp Neurol. 445(3):238-55
- Henderson JM, Annett LE, Torres EM, Dunnett SB (1998) Behavioural effects of subthalamic nucleus lesions in the hemiparkinsonian marmoset (Callithrix jacchus). Eur J Neurosci. 10(2):689-98
- Höhl-Abrahão JC, Creutzfeldt OD (1991) Topographical mapping of the thalamocortical projections in rodents and comparison with that in primates. Exp Brain Res. 87(2):283-94
- Huffman KJ, Krubitzer L (2001) Thalamo-cortical connections of areas 3a and M1 in marmoset monkeys. J Comp Neurol. 435(3):291-310
- 33. Irle E, Markowitsch HJ (1986) Afferent connections of the substantia innominata/basal nucleus of Meynert in car-

- nivores and primates. J Hirnforsch. 27(3):343-67
- 34. Iyengar S, Qi HX, Jain N, Kaas JH (2007) Cortical and thalamic connections of the representations of the teeth and tongue in somatosensory cortex of new world monkeys. J Comp Neurol. 501(1):95-120
- 35. Jacobowitz DM, MacLean PD (1978) A brainstem atlas of catecholaminergic neurons and serotonergic perikarya in a pygmy primate (Cebuella pygmaea). J Comp Neurol. 177(3):397-416
- Kaas JH, Huerta MF, Weber JT, Harting JK (1978) Patterns of retinal terminations and laminar organization of the lateral geniculate nucleus of primates. J Comp Neurol. 182(3):517-53
- 37. Kaske A, Dick A, Creutzfeldt OD (1991) The local domain for divergence of subcortical afferents to the striate and extrastriate visual cortex in the common marmoset (Callithrix jacchus): a multiple labelling study. Exp Brain Res. 84(2):254-65
- Kendrick KM, Dixson AF (1986) Anteromedial hypothalamic lesions block proceptivity but not receptivity in the female common marmoset (Callithrix jacchus). Brain Res. 375(2):221-9
- Kilavik BE, Silveira LC, Kremers J (2003) Centre and surround responses of marmoset lateral geniculate neurones at different temporal frequencies. J Physiol. 546(Pt 3):903-19
- Kilavik BE, Silveira LC, Kremers J (2007) Spatial receptive field properties of lateral geniculate cells in the owl monkey (Aotus azarae) at different contrasts: a comparative study. Eur J Neurosci. 26(4):992-1006
- Kozyrev V, Silveira LC, Kremers J (2007) Linking lateral interactions in flicker perception to lateral geniculate nucleus cell responses. J Physiol. 581(Pt 3):1083-100
- Kremers J (1998-1999) Spatial and temporal response properties of the major retino-geniculate pathways of Old and New World monkeys. Doc Ophthalmol. 95(3-4):229-45
- 43. Kremers J, Kozyrev V, Silveira LC, Kilavik BE (2004) Lateral interactions in the perception of flicker and in the physiology of the lateral geniculate nucleus. J Vis. 4(7):643-63
- 44. Kremers J, Silveira LC, Kilavik BE (2001) Influence of contrast on the responses of marmoset lateral geniculate cells to drifting gratings. J Neurophysiol. 85(1):235-46
- 45. Kremers J, Weiss S (1997) Receptive field dimensions of lateral geniculate cells in the common marmoset (Callithrix jacchus). Vision Res. 37(16):2171-81
- Kremers J, Weiss S, Zrenner E (1997) Temporal properties of marmoset lateral geniculate cells. Vision Res. 37(19):2649-60
- Krubitzer LA, Kaas JH (1992) The somatosensory thalamus of monkeys: cortical connections and a redefinition of nuclei in marmosets. J Comp Neurol. 319(1):123-40
- Leichnetz GR, Astruc J (1975) Efferent connections of the orbitofrontal cortex in the marmoset (Saguinus oedipus). Brain Res. 84(2):169-80
- Lipp HP (1978) Aggression and flight behaviour of the marmoset monkey Callithrix jacchus: an ethogram for brain stimulation studies. Brain Behav Evol. 15(4):241-59
- Lipp HP (1980) A stereotaxic x-ray map of the hypothalamus of the marmoset monkey Callithrix jacchus. Exp Brain Res. 38(2):189-95
- Lipp HP, Hunsperger RW (1978) Threat, attack and flight elicited by electrical stimulation of the ventromedial hypothalamus of the marmoset monkey Callithrix jacchus. Brain Behav Evol. 15(4):260-93
- Markowitsch HJ, Emmans D, Irle E, Streicher M, Preilowski B (1985) Cortical and subcortical afferent connections of the primate's temporal pole: a study of rhesus monkeys, squirrel monkeys, and marmosets. J Comp Neurol. 242(3):425-58
- Markstahler U, Bach M, Spatz WB (1998) Transient molecular visualization of ocular dominance columns (ODCs) in normal adult marmosets despite the desegregated termination of the retino-geniculo-cortical pathways. J Comp Neurol. 393(1):118-34
- Martin PR, White AJ, Goodchild AK, Wilder HD, Sefton AE (1997) Evidence that blue-on cells are part of the third geniculocortical pathway in primates. Eur J Neurosci. 9(7):1536-41
- Pinato L, Allemandi W, Abe LK, Frazão R, Cruz-Rizzolo RJ, Cavalcante JS, Costa MS, Nogueira MI (2007) A
 comparative study of cytoarchitecture and serotonergic afferents in the suprachiasmatic nucleus of primates (Cebus
 apella and Callithrix jacchus) and rats (Wistar and Long Evans strains). Brain Res. 1149:101-10
- Prado Reis F, Abrantes Erhart E (1979) Brain stem, cerebellum and diencephalon of the marmoset (Callithrix jacchus). Acta Anat (Basel). 103(3):278-91
- Qi HX, Lyon DC, Kaas JH (2002) Cortical and thalamic connections of the parietal ventral somatosensory area in marmoset monkeys (Callithrix jacchus). J Comp Neurol. 443(2):168-82
- Ridley RM, Baker HF, Cummings RM, Green ME, Leow-Dyke A (2005) Mild topographical memory impairment following crossed unilateral lesions of the mediodorsal thalamic nucleus and the inferotemporal cortex. Behav Neurosci. 119(2):518-25
- Ridley RM, Baker HF, Mills DA, Green ME, Cummings RM (2004) Topographical memory impairments after unilateral lesions of the anterior thalamus and contralateral inferotemporal cortex. Neuropsychologia. 42(9):1178-91
- Ridley RM, Maclean CJ, Young FM, Baker HF (2002) Learning impairments in monkeys with combined but not separate excitotoxic lesions of the anterior and mediodorsal thalamic nuclei. Brain Res. 950(1-2):39-51
- Schofield SP, Dixson AF (1982) Distribution of catecholamine and indoleamine neurons in the brain of the common marmoset (Callithrix jacchus). J Anat. 134(Pt 2):315-38
- Solomon SG (2002) Striate cortex in dichromatic and trichromatic marmosets: neurochemical compartmentalization and geniculate input. J Comp Neurol. 450(4):366-81

- Solomon SG, White AJ, Martin PR (1999) Temporal contrast sensitivity in the lateral geniculate nucleus of a New World monkey, the marmoset Callithrix jacchus. J Physiol. 517 (Pt 3):907-17
- Solomon SG, White AJ, Martin PR (2002) Extraclassical receptive field properties of parvocellular, magnocellular, and koniocellular cells in the primate lateral geniculate nucleus. J Neurosci. 22(1):338-49
- 65. Spatz WB (1975) Thalamic and other subcortical projections to area MT (visual area of superior temporal sulcus) in the marmoset Callithrix jacchus. Brain Res. 99(1):129-34
- 66. Spatz WB (1978) The retino-geniculo-cortical pathway in Callithrix. I. Intraspecific variations in the lamination pattern of the lateral geniculate nucleus. Exp Brain Res. 33(3-4):551-63
- Spatz WB (1979) The retino-geniculo-cortical pathway in Callithrix. II. The geniculo-cortical projection. Exp Brain Res. 36(3):401-10
- Spatz WB, Erdmann G (1974) Striate cortex projections to the lateral geniculate and other thalamic nuclei: a study using degeneration and autoradiographic tracing methods in the marmoset Callithrix. Brain Res. 82(1):91-108
- Stepniewska I, Ql HX, Kaas JH (2000) Projections of the superior colliculus to subdivisions of the inferior pulvinar in New World and Old World monkeys. Vis Neurosci. 17(4):529-49
- 70. Stevens CF (2002) Predicting functional properties of visual cortex from an evolutionary scaling law. Neuron. 36(1):139-42
- Victor JD, Blessing EM, Forte JD, Buzás P, Martin PR (2007) Response variability of marmoset parvocellular neurons. J Physiol. 579(Pt 1):29-51
- Webb BS, Tinsley CJ, Barraclough NE, Easton A, Parker A, Derrington AM (2002) Feedback from V1 and inhibition from beyond the classical receptive field modulates the responses of neurons in the primate lateral geniculate nucleus, Vis Neurosci. 19(5):583-92
- Webb BS, Tinsley CJ, Vincent CJ, Derrington AM (2005) Spatial distribution of suppressive signals outside the classical receptive field in lateral geniculate nucleus. J Neurophysiol, 94(3):1789-97
- Weiss S, Kremers J, Maurer J (1998) Interaction between rod and cone signals in responses of lateral geniculate neurons in dichromatic marmosets (Callithrix jacchus). Vis Neurosci. 15(5):931-43
- White AJ, Solomon SG, Martin PR (2001) Spatial properties of koniocellular cells in the lateral geniculate nucleus of the marmoset Callithrix jacchus. J Physiol. 533(Pt 2):519-35
- White AJ, Wilder HD, Goodchild AK, Sefton AJ, Martin PR (1998) Segregation of receptive field properties in the lateral geniculate nucleus of a New-World monkey, the marmoset Callithrix jacchus. J Neurophysiol. 80(4):2063-76
- 77. Wilson P, Kitchener PD, Snow PJ (1999) Cutaneous receptive field organization in the ventral posterior nucleus of the thalamus in the common marmoset. J Neurophysiol. 82(4):1865-75
- Yeh T, Lee BB, Kremers J, Cowing JA, Hunt DM, Martin PR, Troy JB (1995) Visual responses in the lateral geniculate nucleus of dichromatic and trichromatic marmosets (Callithrix jacchus). J Neurosci. 15(12):7892-904
- Zhang HQ, Murray GM, Coleman GT, Turman AB, Zhang SP, Rowe MJ (2001) Functional characteristics of the parallel SI- and SII-projecting neurons of the thalamic ventral posterior nucleus in the marmoset. J Neurophysiol. 85(5):1805-22
- Zhang HQ, Murray GM, Turman AB, Mackie PD, Coleman GT, Rowe MJ (1996) Parallel processing in cerebral cortex of the marmoset monkey: effect of reversible SI inactivation on tactile responses in SII. J Neurophysiol, 76(6):3633-55
- Zhang HQ, Zachariah MK, Coleman GT, Rowe MJ (2001) Hierarchical equivalence of somatosensory areas I and II for tactile processing in the cerebral cortex of the marmoset monkey. J Neurophysiol. 85(5):1823-35

Midbrain

- Beckstead RM, Frankfurter A (1982) The distribution and some morphological features of substantia nigra neurons that project to the thalamus, superior colliculus and pedunculopontine nucleus in the monkey. Neuroscience. 7(10):2377-88
- 2. Blanks RH, Clarke RJ, Lui F, Giolli RA, Van Pham S, Torigoe Y (1995) Projections of the lateral terminal accessory optic nucleus of the common marmoset (Callithrix jacchus). J Comp Neurol. 354(4):511-32
- Bourne JA, Rosa MG (2003) Laminar expression of neurofilament protein in the superior colliculus of the marmoset monkey (Callithrix jacchus). Brain Res. 973(1):142-5
- Clarke RJ, Blanks RH, Giolli RA (2003) Midbrain connections of the olivary pretectal nucleus in the marmoset (Callithrix jacchus): implications for the pupil light reflex pathway. Anat Embryol (Berl). 207(2):149-55
- Collins CE, Lyon DC. Kaas JH (2003) Responses of neurons in the middle temporal visual area after long-standing lesions of the primary visual cortex in adult new world monkeys. J Neurosci. 23(6):2251-64
- Collins CE, Lyon DC, Kaas JH (2005) Distribution across cortical areas of neurons projecting to the superior colliculus in new world monkeys. Anat Rec A Discov Mol Cell Evol Biol. 285(1):619-27
- de la Mothe LA, Blumell S, Kajikawa Y, Hackett TA (2006) Thalamic connections of the auditory cortex in marmoset monkeys: core and medial belt regions. J Comp Neurol. 496(1):72-96
- 8. Eslamboli A (2005) Marmoset monkey models of Parkinson's disease: which model, when and why? Brain Res Bull. 68(3):140-9
- Eslamboli A, Romero-Ramos M, Burger C, Bjorklund T, Muzyczka N, Mandel RJ, Baker H, Ridley RM, Kirik D (2007) Long-term consequences of human alpha-synuclein overexpression in the primate ventral midbrain. Brain.


130(Pt 3):799-815


- Everitt BJ, Sirkiä TE, Roberts AC, Jones GH, Robbins TW (1988) Distribution and some projections of cholinergic neurons in the brain of the common marmoset, Callithrix jacchus. J Comp Neurol. 271(4):533-58
- Féger J, Crossman AR (1984) Identification of different subpopulations of neostriatal neurones projecting to globus pallidus or substantia nigra in the monkey: a retrograde fluorescence double-labelling study. Neurosci Lett. 49(1-2):7-12
- 12. Garey LJ, Webster WR (1989) Functional morphology in the inferior colliculus of the marmoset. Hear Res. 38(1-2):67-79
- Hardman CD, Henderson JM, Finkelstein DI, Horne MK, Paxinos G, Halliday GM (2002) Comparison of the basal ganglia in rats, marmosets, macaques, baboons, and humans: volume and neuronal number for the output, internal relay, and striatal modulating nuclei. J Comp Neurol. 445(3):238-55
- Hornung JP, Fritschy JM (1988) Serotoninergic system in the brainstem of the marmoset: a combined immunocytochemical and three-dimensional reconstruction study. J Comp Neurol. 270(4):471-87
- Iczkiewicz J, Jackson MJ, Smith LA, Rose S, Jenner P (2006) Osteopontin expression in substantia nigra in MPTPtreated primates and in Parkinson's disease. Brain Res. 1118(1):239-50
- Jacobowitz DM, MacLean PD (1978) A brainstem atlas of catecholaminergic neurons and serotonergic perikarya in a pygmy primate (Cebuella pygmaea). J Comp Neurol. 177(3):397-416
- Karasawa N, Hayashi M, Yamada K, Nagatsu I, Iwasa M, Takeuchi T, Uematsu M, Watanabe K, Onozuka M (2007)
 Tyrosine hydroxylase (TH)- and aromatic-L-amino acid decarboxylase (AADC)-immunoreactive neurons of the
 common marmoset (Callithrix jacchus) brain: an immunohistochemical analysis. Acta Histochem Cytochem.
 40(3):83-92
- Kirik D, Annett LE, Burger C, Muzyczka N, Mandel RJ, Björklund A (2003) Nigrostriatal alpha-synucleinopathy induced by viral vector-mediated overexpression of human alpha-synuclein: a new primate model of Parkinson's disease. Proc Natl Acad Sci U S A. 100(5):2884-9
- Kunikowska G, Jenner P (2003) Alterations in m-RNA expression for Cu, Zn-superoxide dismutase and glutathione peroxidase in the basal ganglia of MPTP-treated marmosets and patients with Parkinson's disease. Brain Res. 968(2):206-18
- Leichnetz GR, Astruc J (1975) Efferent connections of the orbitofrontal cortex in the marmoset (Saguinus oedipus). Brain Res, 84(2):169-80
- Pow DV, Sullivan RK, Williams SM, Scott HL, Dodd PR, Finkelstein D (2005) Differential expression of the GABA transporters GAT-1 and GAT-3 in brains of rats, cats, monkeys and humans. Cell Tissue Res. 320(3):379-92
- Schofield SP, Dixson AF (1982) Distribution of catecholamine and indoleamine neurons in the brain of the common marmoset (Callithrix jacchus). J Anat. 134(Pt 2):315-38
- Schroeder DM, Jane JA (1976) The intercollicular area of the inferior colliculus. Brain Behav Evol. 13(2-3):125-41
- Sirinathsinghji DJ, Kupsch A, Mayer E, Zivin M, Pufal D, Oertel WH (1992) Cellular localization of tyrosine hydroxylase mRNA and cholecystokinin mRNA-containing cells in the ventral mesencephalon of the common marmoset: effects of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. Brain Res Mol Brain Res. 12(1-3):267-74
- Stepniewska I, Ql HX, Kaas JH (2000) Projections of the superior colliculus to subdivisions of the inferior pulvinar in New World and Old World monkeys. Vis Neurosci. 17(4):529-49
- Svenningsson P, Gunne L, Andren PE (2000) L-DOPA produces strong induction of c-fos messenger RNA in dopamine-denervated cortical and striatal areas of the common marmoset. Neuroscience. 99(3):457-68
- van Vliet SA, Blezer EL, Jongsma MJ, Vanwersch RA, Olivier B, Philippens IH (2008) Exploring the neuroprotective effects of modafinil in a marmoset Parkinson model with immunohistochemistry, magnetic resonance imaging and spectroscopy. Brain Res. 1189:219-28
- vogt Weisenhorn DM, Illing RB, Spatz WB (1995) Morphology and connections of neurons in area 17 projecting to the extrastriate areas MT and 19DM and to the superior colliculus in the monkey Callithrix jacchus. J Comp Neurol. 362(2):233-55
- Waters CM, Hunt SP, Jenner P, Marsden CD (1987) Localization of neurotensin receptors in the forebrain of the common marmoset and the effects of treatment with the neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. Brain Res. 412(2):244-53
- Weber JT, Giolli RA (1986) The medial terminal nucleus of the monkey: evidence for a 'complete' accessory optic system. Brain Res. 365(1):164-8
- 31. Zarate-Lagunes M, Gu WJ, Blanchard V, Francois C, Muriel MP, Mouatt-Prigent A, Bonici B, Parent A, Hartmann A, Yelnik J, Boehme GA, Pradier L, Moussaoui S, Faucheux B, Raisman-Vozari R, Agid Y, Brice A, Hirsch EC (2001) Parkin immunoreactivity in the brain of human and non-human primates: an immunohistochemical analysis in normal conditions and in Parkinsonian syndromes. J Comp Neurol. 432(2):184-96

Stereotaxic Atlas of the Marmoset Brain

with Immunohistochemical Architecture and MR Images

Volume 2
Immunohistochemical
plates and MR images





Shigeki Yuasa Katsuki Nakamura Shinichi Kohsaka

National Institute of Neuroscience National Center of Neurology and Psychiatry, Japan

Stereotaxic Atlas of the Marmoset Brain

with immunohistochemical architecture and MR images

Volume 2

Immunohistochemical plates and MR images

Shigeki Yuasa M.D., Ph.D.

Director, Department of Ultrastructural Research

Katsuki Nakamura Ph.D.

Director, Department of Animal Models for Human Disease

Shinichi Kohsaka M.D., Ph.D.

Director-General

Contents

1.	Gene	eral construction of this atlas	1		
2.	Tables of Plates				
3.	. Plates				
	3.1.	calbindin immunohistochemistry	7		
	3.2.	calretinin immunohistochemistry	69		
	3.3.	tyrosine hydroxylase immunohistochemistry	131		
	3 4	MRI data	175		

1. General construction of this atlas

Contents of Vol. 1

Coronal or parasagittal plates of serial sections stained with cresyl violet (Nissl staining) and the accompanying diagrams with labeling of structures. Corresponding plates of adjacent immunohistochemically-stained sections and MRI data in Vol. 2 are shown in the Tables of plates. Stereotaxic coordinates are indicated in the diagrams.

Contents of Vol. 2

Coronal or parasagittal plates of serial sections immunostained with anti-calbindin, anti-calretinin or anti-tyrosine hydroxylase antibody, and MRI data in the serial coronal or parasagittal sections. Corresponding plates of coronal or parasagittal sections stained with cresyl violet (Nissl staining) and the accompanying diagrams with labeling of structures in Vol. 1 are shown in the Tables of plates.

2. Tables of plates

coronal plane

tracing	Nissl staining	calbindin	calretinin	tyrosine hydroxylase	MRI
cTr-01	cNissl-01				
cTr-02	cNissl-02				MRI cor-0
cTr-03	cNissl-03				
cTr-04	cNissl-04				MRI cor-0
cTr-05	cNissl-05				MRI cor-0
cTr-06	cNissl-06	cCalb-06	cCR-06	cTH-06	MRI cor-0
cTr-07	cNissl-07				
cTr-08	cNissl-08	cCalb-08	cCR-08	cTH-08	
cTr-09	cNissl-09	cCalb-09	cCR-09	cTH-09	MRI cor-0
cTr-10	cNissl-10				
cTr-11	cNissl-11	cCalb-11	cCR-11	cTH-11	
cTr-12	cNissl-12	cCalb-12	cCR-12	cTH-12	MRI cor-0
cTr-13	cNissl-13				
cTr-14	cNissl-14	cCalb-14	cCR-14	cTH-14	
cTr-15	cNissl-15	cCalb-15	cCR-15	cTH-15	MRI cor-0
cTr-16	cNissl-16				MRI cor-0
cTr-17	cNissl-17	cCalb-17	cCR-17	cTH-17	
cTr-18	cNissl-18	cCalb-18	cCR-18	cTH-18	MRI cor-0
cTr-19	cNissl-19	cCalb-19	cCR-19		
cTr-20	(cCalb-20)	cCalb-20	cCR-20		MRI cor-1
cTr-21	cNissl-21	cCalb-21	cCR-21	cTH-21	
cTr-22	cNissl-22	cCalb-22	cCR-22	cTH-22	
cTr-23	cNissl-23				
cTr-24	cNissl-24				MRI cor-1
cTr-25	cNissl-25				
cTr-26	cNissl-26	cCalb-26	cCR-26	cTH-26	
cTr-27	(cCalb-27)	cCalb-27		cTH-27	
cTr-28	(cCalb-28)	cCalb-28	cCR-28	cTH-28	
cTr-29	(cCalb-29)	cCalb-29	cCR-29		MRI cor-1
cTr-30	cNissl-30		cCR-30	cTH-30	
cTr-31	cNissl-31		cCR-31	cTH-31	
cTr-32	(cCalb-32)	cCalb-32	cCR-32	cTH-32	
cTr-33	cNissl-33	cCalb-33	cCR-33	cTH-33	MRI cor-1
cTr-34	cNissl-34	cCalb-34	cCR-34	cTH-34	
cTr-35	cNissl-35	cCalb-35	cCR-35	cTH-35	
cTr-36	cNissl-36	cCalb-36	cCR-36	cTH-36	MRI cor-1
cTr-37	cNissl-37	cCalb-37	cCR-37	cTH-37	
cTr-38	cNissl-38	cCalb-38	cCR-38	cTH-38	
cTr-39	cNissl-39	cCalb-39			MRI cor-1
cTr-40	cNissl-40	cCalb-40		cTH-40	
cTr-41	cNissl-41			cTH-41	
cTr-42	cNissl-42				

coronal plane


tracing	Nissl staining	calbindin	calretinin	tyrosine hydroxylase	MRI
cTr-43	cNissl-43	cCalb-43	cCR-43	cTH-43	MRI cor-16
cTr-44	cNissl-44	cCalb-44	cCR-44	cTH-44	
cTr-45	cNissl-45	cCalb-45	cCR-45	cTH-45	
cTr-46	cNissl-46	cCalb-46	cCR-46	cTH-46	MRI cor-17
cTr-47	cNissl-47	cCalb-47	cCR-47	cTH-47	
cTr-48	cNissl-48	cCalb-48	cCR-48	cTH-48	
cTr-49	(cCalb-49)	cCalb-49	cCR-49	cTH-49	MRI cor-18
cTr-50	cNissl-50	cCalb-50	cCR-50		
cTr-51	cNissl-51	cCalb-51	cCR-51		
cTr-52	cNissl-52	cCalb-52	cCR-52		MRI cor-19
cTr-53	cNissl-53	cCalb-53	cCR-53		MRI cor-20
cTr-54	cNissl-54		cCR-54		MRI cor-21
cTr-55	cNissl-55		cCR-55		
cTr-56	cNissl-56	cCalb-56	cCR-56		MRI cor-22
cTr-57	cNissl-57	cCalb-57	cCR-57		
cTr-58	cNissl-58	cCalb-58	cCR-58		MRI cor-23
cTr-59	cNissl-59	cCalb-59	cCR-59		
cTr-60	cNissl-60	cCalb-60	cCR-60		
cTr-61	cNissl-61	cCalb-61	cCR-61		MRI cor-24
cTr-62	cNissl-62	cCalb-62	cCR-62		
cTr-63	cNissl-63	cCalb-63	cCR-63		MRI cor-25
cTr-64	cNissl-64				MRI cor-26
cTr-65	cNissl-65	cCalb-65	cCR-65		MRI cor-23
cTr-66	cNissl-66	cCalb-66	cCR-66		MRI cor-26

parasagittal plane

tracing	Nissl staining	calbindin	calretinin	tyrosine hydroxylase	MRI
psTr-01	psNissl-01	psCalb-01	psCR-01		MRI psg-01
psTr-02	psNissl-02	psCalb-02	psCR-02	psTH-02	MRI psg-02
psTr-03	psNissl-03	psCalb-03	psCR-03	psTH-03	MRI psg-03
psTr-04	psNissl-04	psCalb-04		psTH-04	
psTr-05	psNissl-05	psCalb-05	psCR-05	psTH-05	MRI psg-04
psTr-06	psNissl-06	psCalb-06	psCR-06	psTH-06	
psTr-07	psNissl-07	psCalb-07	psCR-07	psTH-07	MRI psg-05
psTr-08	psNissl-08	psCalb-08	psCR-08	psTH-08	MRI psg-06
psTr-09	psNissl-09	psCalb-09	psCR-09		MRI psg-07
psTr-10	psNissl-10	psCalb-10	psCR-10	psTH-10	
psTr-11	psNissl-11	psCalb-11	psCR-11	psTH-11	MRI psg-08
psTr-12	psNissl-12	psCalb-12	psCR-12	psTH-12	MRI psg-09
psTr-13	psNissl-13	psCalb-13	psCR-13		MRI psg-10~11
psTr-14	psNissl-14	psCalb-14			MRI psg-12

3. Plates

3.1 Plates—calbindin immunohistochemistry


































































































































3.2 Plates—calretinin immunohistochemistry





































































































































psCR-08




psCR-09





psCR-11





3.3 Plates—tyrosine hydroxylase immunohistochemistry

cTH-06





cTH-08



























































































3.4 Plates—MRI data




21 mm rostral to the interaural line; y=+21

Horizontal line represents the level of the interaural line.

Vertical line represents the level of the interhemispheric fissure.

Scale bar, 10 mm.




18 mm rostral to the interaural line; y=+18


Horizontal line represents the level of the interaural line.

Vertical line represents the level of the interhemispheric fissure.

Scale bar, 10 mm.




16 mm rostral to the interaural line; y=+16
Horizontal line represents the level of the interaural line.
Vertical line represents the level of the interhemispheric fissure.
Scale bar, 10 mm.



15 mm rostral to the interaural line; y=+15

Horizontal line represents the level of the interaural line.


Vertical line represents the level of the interhemispheric fissure.



14 mm rostral to the interaural line; y=+14

Horizontal line represents the level of the interaural line.


Vertical line represents the level of the interhemispheric fissure.




13 mm rostral to the interaural line; y=+13

Horizontal line represents the level of the interaural line.

Vertical line represents the level of the interhemispheric fissure.




12 mm rostral to the interaural line; y=+12
Horizontal line represents the level of the interaural line.
Vertical line represents the level of the interhemispheric fissure.
Scale bar, 10 mm.



11 mm rostral to the interaural line; y=+11


Horizontal line represents the level of the interaural line.

Vertical line represents the level of the interhemispheric fissure.




10 mm rostral to the interaural line; y=+10
Horizontal line represents the level of the interaural line.


Vertical line represents the level of the interhemispheric fissure.



9 mm rostral to the interaural line; y=+9
Horizontal line represents the level of the interaural line.
Vertical line represents the level of the interhemispheric fissure.
Scale bar, 10 mm.




8 mm rostral to the interaural line; y=+8
Horizontal line represents the level of the interaural line.
Vertical line represents the level of the interhemispheric fissure.
Scale bar, 10 mm.




7 mm rostral to the interaural line; y=+7

Horizontal line represents the level of the interaural line.

Vertical line represents the level of the interhemispheric fissure.




6 mm rostral to the interaural line; y=+6
Horizontal line represents the level of the interaural line.
Vertical line represents the level of the interhemispheric fissure.
Scale bar, 10 mm.




5 mm rostral to the interaural line; y=+5

Horizontal line represents the level of the interaural line.

Vertical line represents the level of the interhemispheric fissure.




4 mm rostral to the interaural line; y=+4
Horizontal line represents the level of the interaural line.
Vertical line represents the level of the interhemispheric fissure.
Scale bar, 10 mm.




3 mm rostral to the interaural line; y=+3

Horizontal line represents the level of the interaural line.


Vertical line represents the level of the interhemispheric fissure.



2 mm rostral to the interaural line; y=+2
Horizontal line represents the level of the interaural line.
Vertical line represents the level of the interhemispheric fissure.
Scale bar, 10 mm.




1 mm rostral to the interaural line; y=+1
Horizontal line represents the level of the interaural line.
Vertical line represents the level of the interhemispheric fissure.




0 mm from the interaural line; y=0

Horizontal line represents the level of the interaural line.


Vertical line represents the level of the interhemispheric fissure.



1 mm caudal to the interaural line; y=-1
Horizontal line represents the level of the interaural line.
Vertical line represents the level of the interhemispheric fissure.
Scale bar, 10 mm.




2 mm caudal to the interaural line; y=-2
Horizontal line represents the level of the interaural line.
Vertical line represents the level of the interhemispheric fissure.
Scale bar, 10 mm.



3 mm caudal to the interaural line; y=-3
Horizontal line represents the level of the interaural line.


Vertical line represents the level of the interhemispheric fissure.



4 mm caudal to the interaural line; y=-4

Horizontal line represents the level of the interaural line.


Vertical line represents the level of the interhemispheric fissure.



5 mm caudal to the interaural line; y=-5

Horizontal line represents the level of the interaural line.


Vertical line represents the level of the interhemispheric fissure.




6 mm caudal to the interaural line; y=-6

Horizontal line represents the level of the interaural line.


Vertical line represents the level of the interhemispheric fissure.




7 mm caudal to the interaural line; y=-7
Horizontal line represents the level of the interaural line.
Vertical line represents the level of the interhemispheric fissure.
Scale bar, 10 mm.




0 mm from the interhemispheric fissure; x=0
Horizontal and vertical lines represent the level of the interaural line.
Scale bar, 10 mm.




1.0 mm from the interhemispheric fissure; x=+1Horizontal and vertical lines represent the level of the interaural line.Scale bar, 10 mm.




2.0 mm from the interhemispheric fissure; x=+2
Horizontal and vertical lines represent the level of the interaural line.
Scale bar, 10 mm.




3.0 mm from the interhemispheric fissure; x=+3
Horizontal and vertical lines represent the level of the interaural line.
Scale bar, 10 mm.




4.0 mm from the interhemispheric fissure; x=+4
Horizontal and vertical lines represent the level of the interaural line.
Scale bar, 10 mm.




5.0 mm from the interhemispheric fissure; x=+5
Horizontal and vertical lines represent the level of the interaural line.
Scale bar, 10 mm.



6.0 mm from the interhemispheric fissure; x=+6
Horizontal and vertical lines represent the level of the interaural line.
Scale bar, 10 mm.



7.0 mm from the interhemispheric fissure; x=+7
Horizontal and vertical lines represent the level of the interaural line.
Scale bar, 10 mm.



8.0 mm from the interhemispheric fissure; x=+8
Horizontal and vertical lines represent the level of the interaural line.
Scale bar, 10 mm.



9.0 mm from the interhemispheric fissure; x=+9
Horizontal and vertical lines represent the level of the interaural line.
Scale bar, 10 mm.



10.0 mm from the interhemispheric fissure; x=+10
Horizontal and vertical lines represent the level of the interaural line.
Scale bar, 10 mm.



11.0 mm from the interhemispheric fissure; x=+11
Horizontal and vertical lines represent the level of the interaural line.
Scale bar, 10 mm.